

ROBERT KOCH INSTITUT



Originally published as:

Yvonne Pfeifer, Gottfried Wilharm, Esther Zander, Thomas A. Wichelhaus, Stefan Göttig, Klaus-Peter Hunfeld, Harald Seifert, Wolfgang Witte and Paul G. Higgins
Molecular characterization of *bla*_{NDM-1} in an *Acinetobacter baumannii* strain isolated in Germany in 2007
(2011) Journal of Antimicrobial Chemotherapy, 66 (9), pp. 1998-2001.

DOI: 10.1093/jac/dkr256

This is an author manuscript.

The definitive version is available at: <http://jac.oxfordjournals.org/>

Molecular characterization of *bla*_{NDM-1} in an *Acinetobacter baumannii* strain isolated in Germany in 2007

Yvonne Pfeifer^{1,*}, Gottfried Wilharm¹, Esther Zander², Thomas A. Wichelhaus³, Stefan Göttig³, Klaus-Peter Hunfeld⁴, Harald Seifert², Wolfgang Witte¹ and Paul G. Higgins²

¹Nosocomial Infections, Robert Koch Institute, Burgstraße 37, 38855 Wernigerode, Germany

²Institute for Medical Microbiology, Immunology and Hygiene, University of Cologne, Goldenfelsstrasse 19-21, 50935 Cologne, Germany

³Hospital of Johann Wolfgang Goethe-University Frankfurt/Main, Frankfurt/Main, Germany

⁴Nordwest Medical Center Frankfurt, Frankfurt/Main, Germany

Abstract

Objectives To investigate the genetic environment of the metallo- β -lactamase gene *bla*_{NDM-1} in an *Acinetobacter baumannii* isolated in 2007 in a German hospital.

Methods Antimicrobial susceptibility testing was performed and resistance genes were characterized by PCR amplification and sequencing. Transferability of β -lactam resistance was tested by broth mating assays and transformation of plasmids. The genetic background of *bla*_{NDM-1} was analysed by primer walking. Typing of the *A. baumannii* strain was performed by repetitive extragenic palindromic sequence-based PCR (rep-PCR) using the DiversiLab system.

Results The multidrug-resistant *A. baumannii* isolate harboured β -lactamase genes *bla*_{NDM-1} and intrinsic *bla*_{OXA-64}, but without the insertion sequence ISAb_a1 often located upstream. Transfer of carbapenem resistance by conjugation and transformation failed. Hybridization of isolated plasmid DNA with *bla*_{NDM} probes was not successful. Shotgun cloning of whole genomic DNA and sequence analyses revealed that *bla*_{NDM-1} was located between two insertion elements of ISAb_a125. Furthermore, this *bla*_{NDM-1}-containing transposon structure was integrated into a chromosomal gene encoding a putative *A. baumannii* major facilitator superfamily (MFS) metabolite/H⁺ symporter.

Conclusions The metallo- β -lactamase gene *bla*_{NDM-1} in this *A. baumannii* strain was integrated in the chromosome on a new transposon structure composed of two copies of insertion sequence ISAb_a125. The variability of the genetic environment of *bla*_{NDM-1} likely facilitates the rapid dissemination of this gene within many Gram-negative bacterial species.

Introduction

In recent years an increasing number of reports on the emergence of multidrug-resistant Gram-negative pathogens have been published and in particular it is the emergence of *bla*_{NDM-1} that has created the most concern.^{1,2} Indeed, since its first description in 2009, this metallo- β -lactamase (MBL) has now been found worldwide.^{3,4} NDM-1 has been identified in various Enterobacteriaceae due to localization of *bla*_{NDM-1} on conjugative plasmids, enabling transfer and rapid dissemination of multidrug resistance.⁴ In 2010 *bla*_{NDM-1} and the related *bla*_{NDM-2} gene were also found in *Acinetobacter baumannii*, whereby either plasmid transfer of these genes was confirmed or the genetic location was not further investigated in detail.⁵⁻⁸ Infections with carbapenemase-producing *A. baumannii* are a serious threat because colistin is often the only treatment option. Carbapenem resistance in *A. baumannii* is mainly due to expression of various OXA β -lactamases, such as OXA-23, OXA-58, OXA-40 and OXA-143-related enzymes, as well as overexpression of the intrinsic OXA-51-like enzyme.⁹ Non-OXA-mediated carbapenem resistance is still rare. Here we report the molecular characterization

of the genetic environment of *bla*_{NDM-1} detected in a clinical *A. baumannii* strain from a German hospital.

Materials and methods

Bacterial strains

A. baumannii strain 161/07 was isolated from a patient who had been repatriated to Germany from Serbia in 2007. The case history surrounding this *bla*_{NDM-1}-positive *A. baumannii* strain 161/07 has been described previously.⁶ Standard *A. baumannii* ATCC 19606 and ATCC 17978 were used as recipients for plasmid transformation. Sodium azide-resistant *Escherichia coli* J53 was used as the recipient for transformation and broth mating assays.

Antimicrobial susceptibility

Antimicrobial drug susceptibilities were determined according to the guidelines of the CLSI by broth microdilution (Table S1, available as Supplementary data at [JAC Online](#)) and Etest (bioMérieux, Nürtingen, Germany).¹⁰ The MBL phenotype was confirmed with MBL-Etest and a combined disc diffusion test CDDT (KPC+MBL Confirm ID Kit; Alere GmbH, Cologne, Germany).

Presence of carbapenem resistance genes

Detection of OXA-type carbapenemases and associated insertion elements was performed by PCR and sequence analyses were performed as previously described.^{11,12} Detection of the *bla*_{NDM-1} gene was performed by PCR using primers ndm1_F (5'-CTGAGCACCGCATTAGCC-3') and ndm-1_R (5'-GGCCGTATGAGTGATTGC-3'). For detection of other common *bla*_{OXA} genes (*bla*_{OXA-1,2,9,10}), the following primers were used: oxa1_F (5'-TATCTACAGCAGCGCCAGTG-3'); oxa1_R (5'-TAAATTCGACCCCAAGTTTCC-3'); oxa2_F (5'-GCAAGAAGGCACGCTAGAAC-3'); oxa2_R (5'-CAGCGTCCGAGTTGACTG-3'); oxa9_F (5'-TTTGCTGCTGCATATGTTGG-3'); oxa9_R (5'-CCCATCAACACGGGTAATTC-3'); oxa10_F (5'-TTCGAGTACGGCATTAGCTG-3'); and oxa10_R (5'-CAATGATGCCCTCACTTTCC-3').

Molecular typing

Strain typing was performed by rep-PCR using the DiversiLab system (bioMérieux) and results were compared with our database of worldwide clonal lineages.⁹ In addition, multiplex PCR-based typing in combination with Apal PFGE was performed.¹³

Characterization of *bla*_{NDM-1}

Transfer of β -lactam resistance was tested by broth mating assays with *E. coli* J53 as the recipient. Selection of transconjugants was performed on Mueller–Hinton agar plates that contained sodium azide (200 mg/L) and ampicillin (100 mg/L). Plasmid DNA was isolated using the QIAGEN Plasmid Mini Kit (QIAGEN, Hilden, Germany).

Transformation of plasmids into an electrocompetent *E. coli* J53 and *A. baumannii* ATCC 19606 and ATCC 17978 recipients was performed using standard procedures.¹⁴ Plasmid size was determined by performing S1 nuclease restriction of whole genomic DNA combined with PFGE.¹⁵ Southern hybridization using digoxigenin-dUTP-labelled probes and signal detection using CDP-*Star* were performed following the manufacturer's guidelines (Roche Diagnostics Ltd, West Sussex, UK).

Whole genomic DNA was isolated using the QIAGEN DNeasy kit and digested with EcoRV, ligated into EcoRV-cut plasmid pBBR1MCS, transformed into *E. coli* NEB 5-alpha and selected on ticarcillin

(25 mg/L).¹⁶ Inserts of isolates that grew on ticarcillin were amplified by PCR using M13 primers and sequenced. Insert DNA was sequenced on both strands by primer walking.

Additionally, transfer of naked whole genomic DNA of *A. baumannii* 161/07 into an ampicillin-susceptible *A. baumannii* 102/07 recipient by natural transformation was performed. We have recently identified conditions promoting the uptake of naked DNA by a significant number of *A. baumannii* isolates. These results will be published elsewhere (E. Skiebe and G. Wilharm, unpublished results).

Nucleotide sequence accession number

The nucleotide and protein sequences of the *bla*_{NDM-1}-containing transposon and gene *aphA-6* have been registered in GenBank under accession numbers HQ857107 and JF949760, respectively.

Results and discussion

Strain typing by rep-PCR (DiversiLab) revealed that the NDM-1-producing multidrug-resistant *A. baumannii* strain 161/07 clustered with isolates of the clonal lineage WW7. Previous work has shown that *A. baumannii* strains belonging to the WW7 cluster harboured the carbapenemase genes *bla*_{OXA-58} or *bla*_{OXA-23} and were from different countries in South America, Europe and Asia.⁹ Furthermore, all *A. baumannii* of the WW7 cluster harboured *bla*_{OXA-64}, a variant of the intrinsic *bla*_{OXA-51}.¹⁷ It was confirmed by PCR that *A. baumannii* 161/07 was positive for the chromosomally located *bla*_{OXA-64}, but this gene was not associated with insertion element *ISAbal*. Other *bla*_{OXA} genes were not detected. PCR-based *A. baumannii* typing in combination with *Apal* PFGE analysis confirmed that *A. baumannii* 161/07 was not related to European clonal lineages 1–3.¹³

Transfer of resistance genes by *in vitro* conjugation and transformation of plasmids into *E. coli* or *A. baumannii* recipients was not successful. Analysis of S1-digested DNA fragments revealed the presence of two plasmids (125 kb, 75 kb) in the NDM-1-producing *A. baumannii* 161/07 strain. However, hybridization signals with a *bla*_{NDM-1} probe were not detected for these plasmids, and PCR with *bla*_{NDM-1}-specific primers failed to amplify a PCR product, suggesting a chromosomal location of *bla*_{NDM-1}.

Using naked whole genomic DNA from *A. baumannii* 161/07 to transform the naturally competent ampicillin-susceptible *A. baumannii* recipient 102/07, the gene *bla*_{NDM-1} was successfully transferred, as confirmed by PCR. All β -lactams tested against the transformant 161/07-102 showed an increase in MIC, with imipenem and meropenem MICs increasing from 0.25 mg/L to >32 mg/L, and 0.5 mg/L to 32 mg/L, respectively (Table S1). No other class of antibiotic was affected.

Sequence analysis of shotgun cloned *A. baumannii* 161/07 genomic DNA revealed a 3.9 kb insert containing the *bla*_{NDM-1} gene and parts of the plasmid sequence pKpANDM-1 described previously (Figure 1C).³ However, the IS26 transposase upstream of *bla*_{NDM-1} in previously characterized plasmids pKpANDM-1 and pNDM-HK was not present in *A. baumannii* 161/07. Instead, the insertion sequence *ISAb125* was identified (Figure 1B). Interestingly, pKpANDM-1 includes a partial sequence of *ISAb125* adjacent to *bla*_{NDM-1}. BLAST analysis of *ISAb125* revealed seven copies of *ISAb125* on the chromosome of *A. baumannii* strain ACICU in the GenBank database, and *ISAb125* was found disrupting the *carO* gene, leading to carbapenem resistance.¹⁸ Furthermore, *ISAb125* can also be plasmid located, and has been described recently upstream of the *bla*_{OXA-58} gene.¹⁹

To determine whether *bla*_{NDM-1} was located in a transposon, PCR was performed with inverse primers to *ISAb125* [TRANSIS_R (5'-AAACAACGGATCGCTTCAAC-3') and TRANSIS_F (5'-CGAGCATTACCAAAGGGTGA-3')] using genomic DNA of *A. baumannii* 161/07 as template. Two products of 2 kb and 9 kb were amplified. Sequencing of the 2 kb amplicon revealed *aphA-6*, an aminoglycoside resistance gene bracketed between two copies of *ISAb125*.

Primer walking from both ends of the 9 kb amplicon revealed a composite transposon structure containing *bla*_{NDM-1} bracketed between two copies of *ISAb125* (Figure 1B). Both copies of *ISAb125*

were flanked by 17 bp inverted repeats. Both 5' inverted repeats and both 3' inverted repeats were identical, respectively, but there were two nucleotide differences between 5' and 3' inverted repeats. The two IS*Aba125* transposase genes differed by four nucleotides whereby one resulted in amino acid substitution, Arg-41→Gln. Including both IS*Aba125* insertion elements, the composite transposon was 10093 bp in length. Furthermore, it was integrated into a chromosomal gene encoding a putative *A. baumannii* major facilitator superfamily (MFS) metabolite/H⁺ symporter (Figure 1A) that has previously been disrupted by IS*Aba1-bla_{OXA-23}*.²⁰ Further PCR and sequencing with primers for IS*Aba125* and the disrupted MFS transporter confirmed the chromosomal location of the *bla_{NDM-1}*-containing transposon. Evidence of this being a transposition event was an 8 bp target site duplication at the point of insertion in the gene encoding MFS (Figure 1E). Adjacent to the MFS gene we identified a chromosomal homoserine lactone synthase gene (*cepI*). In addition, inside the *bla_{NDM-1}*-containing transposon there was a 4 kb element 93% similar to that described in *E. coli* plasmids pEH4H and pAR060302, encoding the chaperonin subunits *groS* and *groL*, and the transposase *insE*, which was not associated with inverted repeats (Figure 1D). The 8 bp target site duplication was also present at the 5'-end of the *insE* gene (Figure 1E). Further BLAST analysis revealed a 297 bp sequence, including 41 bp of the 3'-end of *insE* and extending towards IS*Aba125*, that showed 99% identity to insertion sequence IS*SCR19*-like and *oriIS* previously described in a *Pseudomonas aeruginosa* isolate.²¹ Deletions were found in gene *trpF* (79 bp, 3'-end) and gene *groS* (154 bp, 5'-end). No significant DNA homology or open reading frames were detected in the 1.6 kb region between the truncated *trpF* and *groS* genes.

In conclusion, analysis of the genetic environment of *bla_{NDM-1}* in *A. baumannii* 161/07 revealed a transposon structure composed of two copies of insertion sequence IS*Aba125* that is integrated into the bacterial chromosome. However, since *bla_{NDM-1}* is flanked by these insertion elements, we cannot discount the possibility of integration into a plasmid and subsequent horizontal spread. The variability of the genetic environment of *bla_{NDM-1}*, as evidenced by the transposon structure described here, with DNA seemingly originating from Enterobacteriaceae, *P. aeruginosa* and *A. baumannii*, may explain the observed rapid dissemination of this gene within many Gram-negative bacterial species and across genera. Thus there is an urgent need for further investigations to find the origin of this gene and its mechanisms of spread.

Funding

This work was funded by the Ministry of Health, Germany. The contribution of H. S. and P. G. H. was supported by a grant from Bundesministerium für Bildung und Forschung (BMBF), Germany, Klinische Forschergruppe Infektiologie (grant number 01 KI 0771).

Transparency declarations

None to declare.

Supplementary data

Table S1 is available as Supplementary data at [JAC Online](http://jac.oxfordjournals.org/) (<http://jac.oxfordjournals.org/>).

Acknowledgements

We thank George A. Jacoby for providing the *E. coli* J53 Azi^r strain. We extend special thanks to Sibylle Müller-Bertling for performing phenotypical and genotypical analyses and Evelyn Skiebe for performing transformation experiments.

References

- 1 Bonomo RA. New Delhi metallo- β -lactamase and multidrug resistance: a global SOS? *Clin Infect Dis* 2011; 52: 485–7.
- 2 Nordmann P, Poirel L, Toleman MA et al. Does broad-spectrum β -lactam resistance due to NDM-1 herald the end of the antibiotic era for treatment of infections caused by Gram-negative bacteria? *J Antimicrob Chemother* 2011; 66: 689–92.
- 3 Yong D, Toleman MA, Giske CG et al. Characterization of a new metallo- β -lactamase gene, blaNDM-1, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 2009; 53: 5046–54.
- 4 Kumarasamy KK, Toleman MA, Walsh TR et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis* 2010; 10: 597–602.
- 5 Karthikeyan K, Thirunarayan MA, Krishnan P. Coexistence of blaOXA-23 with blaNDM-1 and armA in clinical isolates of *Acinetobacter baumannii* from India. *J Antimicrob Chemother* 2010; 65: 2253–4.
- 6 Goettig S, Pfeifer Y, Wichelhaus TA et al. Global spread of New Delhi metallo- β -lactamase 1. *Lancet Infect Dis* 2010; 10: 828–9.
- 7 Chen Y, Zhou Z, Jiang Y et al. Emergence of NDM-1-producing *Acinetobacter baumannii* in China. *J Antimicrob Chemother* 2011; 66: 1255–9.
- 8 Kaase M, Nordmann P, Wichelhaus TA et al. NDM-2 carbapenemase in *Acinetobacter baumannii* from Egypt. *J Antimicrob Chemother* 2011; 66: 1260–2.
- 9 Higgins PG, Dammhayn C, Hackel M et al. Global spread of carbapenem-resistant *Acinetobacter baumannii*. *J Antimicrob Chemother* 2010; 65: 233–8.
- 10 Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing: Twenty-first Informational Supplement M100-S21. CLSI, Wayne, PA, USA, 2011.
- 11 Woodford N, Ellington MJ, Coelho JM et al. Multiplex PCR for genes encoding prevalent OXA carbapenemases in *Acinetobacter* spp. *Int J Antimicrob Agents* 2006; 27: 351–3.
- 12 Turton JF, Ward ME, Woodford N et al. The role of ISAbal in expression of OXA carbapenemase genes in *Acinetobacter baumannii*. *FEMS Microbiol Lett* 2006; 258: 72–7.
- 13 Turton JF, Gabriel SN, Valderrey C et al. Use of sequence-based typing and multiplex PCR to identify clonal lineages of outbreak strains of *Acinetobacter baumannii*. *Clin Microbiol Infect* 2007; 13: 807–15.
- 14 Choi KH, Kumar A, Schweizer HP. A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Methods* 2006; 64: 391–7.
- 15 Barton BM, Harding GP, Zuccarelli AJ. A general-method for detecting and sizing large plasmids. *Anal Biochem* 1995; 226: 235–40.
- 16 Kovach ME, Phillips RW, Elzer PH et al. pBBR1MCS: a broad-host-range cloning vector. *Biotechniques* 1994; 16: 800–2.
- 17 Zander E, Higgins PG, Seifert H. Correlation between blaOXA-51 variant and DiversiLab worldwide clonal lineages in *Acinetobacter baumannii*. In: Abstracts of the Twenty-first European Congress of Clinical Microbiology and Infectious Diseases/Twenty-seventh International Congress of Chemotherapy, Milan, Italy, 2011. Abstract P1731. European Society of Clinical Microbiology and Infectious Diseases, Basel, Switzerland.
- 18 Mussi MA, Limansky AS, Viale AM. Acquisition of resistance to carbapenems in multidrug-resistant clinical strains of *Acinetobacter baumannii*: natural insertional inactivation of a gene encoding a member of a novel family of β -barrel outer membrane proteins. *Antimicrob Agents Chemother* 2005; 49: 1432–40.
- 19 Evans BA, Hamouda A, Towner KJ et al. Novel genetic context of multiple blaOXA-58 genes in *Acinetobacter* genospecies 3. *J Antimicrob Chemother* 2010; 65: 1586–8.
- 20 Meric M, Kasap M, Gacar G et al. Emergence and spread of carbapenem-resistant *Acinetobacter baumannii* in a tertiary care hospital in Turkey. *FEMS Microbiol Lett* 2008; 282: 214–8.
- 21 Naas T, Namdari F, Bogaerts P et al. Genetic structure associated with blaOXA-18, encoding a clavulanic acid-inhibited extended-spectrum oxacillinase. *Antimicrob Agents Chemother* 2008; 52: 3898–904.

Figure

Figure 1. Schematic diagram showing the genetic environment of *bla_{NDM-1}* in *A. baumannii* 161/07. (A) Chromosomal genes *cepl* and the truncated *mfs*; (B) *ISAbal25*; (C) *bla_{NDM-1}* and truncated *trpF*; (D) truncated chaperonin subunit *groS*, chaperonin *groL* and the transposase *insE*; (E) 8 bp target duplication.

