



Weekly report of 20-04-2022

Reporting period: 24-06-2021 – 17-04-2022

ABOUT THE WEEKLY REPORT

SUMO is a system that has been developed and implemented at the Robert Koch Institute. It processes and provides health data for surveillance and public health research. The emergency department weekly report presents data from the AKTIN emergency department registry from the routine documentation of selected German emergency departments. The report shows the current utilisation of these emergency departments as part of the emergency department surveillance at the Robert Koch Institute.

Data source and inclusion criteria

The inclusion of emergency departments is based on voluntary participation and is determined per quarter, depending on the data availability.

Emergency departments are included in the report, if data gaps where no emergency department admissions are reported are not larger than 6 consecutive days each. Furthermore, to be included in the most recent quarter an emergency department has to report at least one admission for each of the last 7 days. These criteria equally apply to the analyses of triage and chief complaints (according to CEDIS-PCL), but consider the availability of the respective variable. To be included in the analyses for syndromic surveillance, an emergency department has to report at least one case of the respective case definition in the quarter.

Because the availability of data can vary by variable and change over time, the number of emergency departments can vary between quarters, between reports and between the different plots. Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

Variables and case definitions

Emergency departments document triage levels based on the *Emergency Severity Index* (ESI) or the *Manchester Triage System* (MTS). The reported chief complaints were coded according to the *Canadian Emergency Department Information System – Presenting Complaint List* (CEDIS-PCL) and diagnoses according to the *International Classification of Diseases – Version 10* (ICD-10).

Case definitions for acute respiratory illness (ARI) and severe ARI (SARI) were based on a combination of diagnosis, chief complaints and hospitalisation, as described in Boender et al. 2021 (DOI 10.1101/2021.08.19.21262303). The case definition for influenza-like illness (ILI) was based on ICD-10 diagnoses: J09, J10.-, J11.-. Case definitions for respiratory, cardiovascular and neurological complaints were based on CEDIS-PCL codes: RC – respiratory (651-661), CV – cardiovascular (001-012) and NC – neurological (401-411).

Baseline calculation

For each category (visits, stratified by age and triage, case definitions) we calculate a baseline and a 95% prediction interval per single emergency department, which is then summed up over all emergency departments to one final baseline and prediction interval, respectively. A negative binomial regression model is used to calculate the baseline. We use a global regression model that is trained on all available emergency departments, allowing the inclusion of emergency departments without full historic data (from 2017-01-01 to 2022-03-31). The regression model includes variables for the respective emergency department, seasonality, a time trend, and pandemic phases. From that we can calculate a pre-pandemic baseline for all emergency departments and points in time. The sum over all included emergency departments yields the depicted total baseline.

Interpretation of the data




A meaningful interpretation of the data is only possible with knowledge of the processes and structures of the respective emergency departments and after consulting with these. Furthermore, for the interpretation of surveillance data, it is important to consider several limitations:

- The sample of emergency departments is not representative for Germany.
- The number of emergency departments in the current quarter may change weekly, because not every emergency department delivers data every week.
- Population based data such as incidences cannot be derived from the presented numbers.
- Complaints are not equal to clinically confirmed diagnoses. Also diagnoses made in emergency departments are often suspected diagnoses.
- The indicators for the syndromic surveillance are defined by a combination of complaints and ICD-10 diagnoses both of which may have missing values (see Table 1). This can lead to an underestimation of case numbers.
- Changes over time can be caused both by real changes of the emergency department utilisation, as well as several other reasons (e.g. changed documentation practices or care processes). Therefore, the data should not be interpreted without prior direct communication with the emergency departments.

PARTNERSHIP


The report has been established in close cooperation with the AKTIN Emergency Department Data Registry. We want to especially thank the participating emergency departments for sharing their data.

**CONTACT****Robert Koch Institute**

 SUMO@rki.de
 www.rki.de/sumo
 Robert Koch Institute
 Nordufer 20
 13353 Berlin, Germany

AKTIN German Emergency Department Data Registry

 office@aktin.org
 www.aktin.org/en
 University Clinic for Traumatology,
 Medical Faculty
 Otto von Guericke University Hospital
 Magdeburg
 Leipziger Straße 44
 39120 Magdeburg, Germany

 Department of Medical Informatics
 Uniklinik RWTH Aachen
 Pauwelsstraße 30
 52074 Aachen

Archive

Previous reports can be found here: [Archive emergency department situation report.](#)

Suggested citation:

Robert Koch Institute: Routine health data in real-time (SUMO). Emergency Department Surveillance Weekly Report 20-04-2022. DOI 10.25646/9831.

WEEKLY OVERVIEW

Reporting date: 20-04-2022
Reporting period: 24-06-2021 – 17-04-2022

Emergency departments: in total 22 emergency departments in Germany, located in the federal states Baden-Wuerttemberg, Bavaria, Lower Saxony, North Rhine-Westphalia, Saxony, Saxony-Anhalt, and Schleswig-Holstein

Level of care:

Basic emergency care: 1 departments
 Extended emergency care: 3 departments
 Comprehensive emergency care: 18 departments




Figure 1: Emergency departments included in the current quarter (green) and included in the previous quarters in this report (grey).

Data quality


Date	Number of attendances	Average number of attendances per department	Triage available (%)	CEDIS-PCL available (%)	ICD-10 diagnosis available (%)	Disposition available (%)
17-04-2022	2,014	91	94.9	64.2	24.8	74.5
16-04-2022	2,264	102	93.9	63.3	29.6	76.4
15-04-2022	2,150	97	95.3	62.7	32.5	74.5
14-04-2022	2,313	105	95.1	62.6	35.5	76.5
13-04-2022	2,279	103	94.0	64.6	49.9	77.8
12-04-2022	2,225	101	94.7	68.4	54.3	77.9
11-04-2022	2,503	113	95.5	65.5	54.4	78.7

Table 1: Number of recorded emergency department attendances and availability of variables for each day of the current week

EMERGENCY DEPARTMENT ATTENDANCES

Figure 2: Emergency department attendances

Daily emergency department attendances shown as 7-day moving average compared to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

AGE

Figure 3a: Overview of all age groups

7-day moving average of emergency department attendances stratified by age groups.





Figure 3b: Attendances per age group compared to baseline

7-day moving average of emergency department attendances stratified by age groups in comparison to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

TRIAGE

Figure 4a: Overview of all triage levels

7-day moving average of emergency department attendances stratified by triage levels.





Figure 4b: Attendances per triage level compared to baseline

7-day moving average of emergency department attendances stratified by triage levels in comparison to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

SYNDROMIC SURVEILLANCE

Figure 5: Acute respiratory infection (ARI)

Daily emergency department attendances for acute respiratory infection (ARI) and 7-day moving average in comparison to baseline.





Figure 6: Severe acute respiratory infection (SARI)

Daily emergency department attendances for severe acute respiratory infection (SARI) and 7-day moving average in comparison to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

Figure 7: Influenza-like-illness (ILI)

Daily emergency department attendances for influenza-like illness (ILI) and 7-day moving average in comparison to baseline.





Figure 8: Respiratory complaints

Daily emergency department attendances for respiratory complaints and 7-day moving average in comparison to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.

Figure 9: Cardiovascular complaints

Daily emergency department attendances for cardiovascular complaints and 7-day moving average in comparison to baseline.





Figure 10: Neurological complaints

Daily emergency department attendances for neurological complaints and 7-day moving average in comparison to baseline.



Note: Because the size of the included emergency departments varies, the number of reported cases can vary between quarters with the same number of emergency departments.