

Frequent Combination of Antimicrobial Multiresistance and Extraintestinal Pathogenicity in *Escherichia coli* Isolates from Urban Rats (*Rattus norvegicus*) in Berlin, Germany

Sebastian Guenther^{1*}, Astrid Bethe¹, Angelika Fruth², Torsten Semmler¹, Rainer G. Ulrich³, Lothar H. Wieler¹, Christa Ewers^{1,4}

1 Freie Universität Berlin, Department of Veterinary Medicine, Institute of Microbiology and Epizootics, Berlin, Germany, **2** Robert Koch-Institut, Wernigerode, Germany, **3** Friedrich-Loeffler-Institut, Federal Research Institute for Animal Health, Institute for Novel and Emerging Infectious Diseases, Greifswald - Insel Riems, Germany, **4** Justus-Liebig-Universität Giessen, Department of Veterinary Medicine, Institute of Hygiene and Infectious Diseases of Animals, Giessen, Germany

Abstract

Urban rats present a global public health concern as they are considered a reservoir and vector of zoonotic pathogens, including *Escherichia coli*. In view of the increasing emergence of antimicrobial resistant *E. coli* strains and the on-going discussion about environmental reservoirs, we intended to analyse whether urban rats might be a potential source of putatively zoonotic *E. coli* combining resistance and virulence. For that, we took fecal samples from 87 brown rats (*Rattus norvegicus*) and tested at least three *E. coli* colonies from each animal. Thirty two of these *E. coli* strains were pre-selected from a total of 211 non-duplicate isolates based on their phenotypic resistance to at least three antimicrobial classes, thus fulfilling the definition of multiresistance. As determined by multilocus sequence typing (MLST), these 32 strains belonged to 24 different sequence types (STs), indicating a high phylogenetic diversity. We identified STs, which frequently occur among extraintestinal pathogenic *E. coli* (ExPEC), such as STs 95, 131, 70, 428, and 127. Also, the detection of a number of typical virulence genes confirmed that the rats tested carried ExPEC-like strains. In particular, the finding of an Extended-spectrum beta-lactamase (ESBL)-producing strain which belongs to a highly virulent, so far mainly human- and avian-restricted ExPEC lineage (ST95), which expresses a serogroup linked with invasive strains (O18:NM:K1), and finally, which produces an ESBL-type frequently identified among human strains (CTX-M-9), pointed towards the important role, urban rats might play in the transmission of multiresistant and virulent *E. coli* strains. Indeed, using a chicken infection model, this strain showed a high *in vivo* pathogenicity. Imagining the high numbers of urban rats living worldwide, the way to the transmission of putatively zoonotic, multiresistant, and virulent strains might not be far ahead. The unforeseeable consequences of such an emerging public health threat need careful consideration in the future.

Citation: Guenther S, Bethe A, Fruth A, Semmler T, Ulrich RG, et al. (2012) Frequent Combination of Antimicrobial Multiresistance and Extraintestinal Pathogenicity in *Escherichia coli* Isolates from Urban Rats (*Rattus norvegicus*) in Berlin, Germany. PLoS ONE 7(11): e50331. doi:10.1371/journal.pone.0050331

Editor: Stephen V. Gordon, University College Dublin, Ireland

Received: June 19, 2012; **Accepted:** October 17, 2012; **Published:** November 26, 2012

Copyright: © 2012 Guenther et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Federal Ministry of Education (BMBF) and Research Network Zoonosis (FBI-Zoo, Grant 01K1012A), the German Research Foundation (DFG) funded Indo-German Research Training Group (Grant GRK1673), and the Federal Ministry of Education and Research grant to RGU (FKZ 01K1018). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sebastian.guenther@fu-berlin.de

Introduction

Brown rats (*Rattus norvegicus*) are commensal rodents found in urban areas worldwide. They are associated with hygienic problems and are considered a reservoir and vector of several zoonotic pathogens. Indeed, until the twentieth century, one of the most feared diseases related to rats was the plague caused by *Yersinia pestis* [1,2]. Nowadays, a number of other bacterial, viral and parasitic pathogens have been associated with rats, such as *Leptospira* spp., Shiga toxin producing *E. coli*, *Campylobacter* spp., *Salmonella* spp., or Hantaviruses [1,3,4,5,6]. There are numerous ways, by which rodent-borne pathogens may infect human and animal hosts. Inhalation of aerosols and consumption of contaminated food are considered the main pathways, while also direct contact, e.g. by bites, or infections via vectors might occur. Even surface water contaminated with droppings and urine from

infected rats in recreational areas has been identified as possible infection source [7]. In addition, specific ecological and behavioral characteristics, e.g. a concentration of Brown rats into high-density populations along with their cohabitation with humans, may further promote the spread of zoonotic pathogens [7].

Another aspect of Brown rats synanthropism is their inhabitation of areas near anthropogenically created food sources, such as garbage or sewage systems, also providing harborage [8,9]. Although it is well known that rats live in certain parts of the sewage system [10], even continuous baiting programs have failed to eliminate Brown rat populations [11]. Brown rats from rural areas can roam as far as 260–2000 m within a day, while observational studies in city environments identified smaller activity areas of 25–150 m for rats in urban areas [10]. Nevertheless, urban Brown rats also appear to be able to build

an epidemiological bridge between the sewage system and populated urban environments, as social factors, such as aggression in case of overpopulation of rats [12] or large disturbances in their environment can force populations to travel long distances also [13]. This can lead to large population fluctuations and the transmission of pathogens hosted by rats into new areas [14].

Although a natural fear of wild rats as putative carriers of infectious agents is largely embedded in our culture [15], there are hardly any scientific data regarding actual population trends. Estimations about the number of animals are scant or not available at all, like is also the case for our study site, Berlin. For other comparable urban areas, the total number of Brown rats seems to have been on a continuous high level over the last 50 years, as it has been reported for Baltimore (USA) [7]. But in recent years, there have also been reports on increased levels of infestation of urban areas in Great Britain [16]. At the same time, there is evidence of substantial under-reporting of rat infestations [10]. Furthermore, a deteriorating integrity of sewage infrastructures combined with less sewer baiting programs [10] may have intensified the occasion of direct and indirect contact between rat and humans in an urban environment. On a global level, climate change and changing human settlement patterns like the ongoing urbanization trend could lead to increased problems with rat-borne pathogens as the distribution of rodent species and pathogens linked to these species could be influenced [10].

Rats are natural hosts of *Escherichia (E.) coli*, a commensal ubiquitous bacterium colonizing the gut of mammals and birds [17]. Here, from a zoonotic perspective, intestinal pathogenic subtypes of *E. coli* (InPEC), including Shiga toxin producing *E. coli* (STEC), enterohemorrhagic *E. coli* (EHEC), enteroaggregative *E. coli* (EaggEC), and extraintestinal pathogenic *E. coli* (ExPEC) are of major concern. Recent studies on the occurrence of putatively zoonotic *E. coli* in rats were largely focused on STEC and on the epidemiologic relevance that rats, living on or in close proximity to cattle farms, might play in the distribution of EHEC O157 isolates [3,5,18]. Yet barely anything is known about the role of rats as carriers of ExPEC, which have received large attention recent years as they often express a multiresistance phenotype. In particular Extended-spectrum beta-lactamase (ESBL)-producing ExPEC strains account for serious problems in the treatment of infectious diseases in humans and animals as these enzymes confer resistance to nearly all beta-lactam antimicrobial drugs, including third-generation cephalosporins [19,20,21,22,23]. Although ESBL strains have been observed among all phylogenetic groups of *E. coli* including “extraintestinal pathogenic groups” B2 and D, still a larger proportion belongs to phylogenetic lineages and multilocus sequence types (STs) that are composed of opportunistic pathogens and commensals, lacking an extensive set of virulence-associated genes and causing infections primarily in immuno-compromised hosts [19,21]. A well-known exception is the worldwide emerging clonal group O25b:H4-B2-ST131-CTX-M-15 which has been implicated in a wide range of severe hospital- and community-acquired extraintestinal infections in humans and animals [24]. The recent finding of an ESBL-producing *E. coli* strain belonging to this pandemic group in the feces of an urban rat from Berlin [22] prompted us to screen urban rats also for other multiresistant *E. coli*, the presence of genes associated with extraintestinal pathogenic and Shiga toxin producing strains, and their phylogenetic relatedness to human and animal clinical strains, determined by multi locus sequence typing (MLST). As the simple possession of virulence associated genes does not necessarily translate to *in vivo* pathogenicity, we chose one exemplary isolate to assess its pathogenicity in a chicken infection model. The strain was selected as it harbored a frequently encountered ESBL type (CTX-M-9)

and represented a prominent and highly invasive ExPEC-lineage (ST95), which so far has been particularly associated with pathogenic human and avian strains and only scarcely expressed a multiresistance phenotype. The data obtained here might help to gain further insight into the role of synanthropic rodents as carriers, reservoir and even disseminators of *E. coli* that combine multiresistance and extraintestinal virulence.

Materials and Methods

Ethics Statement

All animal experiments were approved by the “Landesamt fuer Gesundheit und Soziales” (Reg. 0220/06) and chickens were killed according to animal welfare norms.

Bacterial Strains

Fecal samples of 87 urban brown rats (*Rattus norvegicus*) were collected on 53 different sampling locations all over Berlin (Germany) from 2008–2009. Rats were captured and euthanized by pest control technicians during pest control (n = 40), or swabs were taken directly at the place of capturing and transferred into conservation medium (Mast Diagnostics, Reinfeld, Germany) (n = 47). After overnight cultivation on ChromeOrientation® agar (Mast Diagnostics) at 37°C, at least three *E. coli* isolates were obtained from each fecal sample. Classical biochemical methods were used to determine the bacterial species [25]. Copy clones recovered from individual animals (copy clones among different individuals were not detected), were excluded by randomly amplified polymorphic DNA (RAPD)-PCR, performed as recently described [26].

Determination of Phenotypic Resistance and Pre-selection of Strains

Preliminary screening for antimicrobial resistance was done by agar dilution test with six different antimicrobial substances as recently described [27]. Here, freshly prepared Mueller–Hinton-agarose plates containing estimated breakpoint concentrations of ampicillin (≥ 32 µg/ml), streptomycin (≥ 64 µg/ml), spectinomycin (≥ 128 µg/ml), chloramphenicol (≥ 32 µg/ml), gentamicin (≥ 16 µg/ml) and tetracycline (≥ 16 µg/ml) were used. Isolates displaying phenotypic resistance for at least one antimicrobial class were additionally tested by Agar broth microdilution method (Micronaut breakpoint plate “Kleintier”, Genzyme Diagnostics, Rüsselsheim, Germany) against seventeen antimicrobials including beta-lactams as well as non-beta-lactams like aminoglycosides, tetracyclines, sulfonamides, chloramphenicol and fluoroquinolones according to the standards given by the CLSI guideline [28]. Phenotypic screening for ESBL production was performed using the confirmatory test with cefotaxime and ceftazidime alone or in combination with clavulanic acid according to the method recommended in the CLSI document M31-A3 [28].

Determination of Antimicrobial Resistance Genes

Multiresistant *E. coli* isolates were screened for the presence of antimicrobial resistance genes, such as *tet(A-D)*, *sul1*, *sul2*, *sul3*, *strA*, *strB*, *aadA1-like*, *aac(3)-IV*, *bla_{TEM-1-like}*, *bla_{SHV}* and *bla_{CTX-M}* using standard PCR methods and sequencing of the PCR products if necessary. The presence of plasmid-mediated quinolone resistance gene variant *aac(6′)-Ib-cr* and the *qnrA*, *qnrB*, and *qnrS* genes as well as of mutations in *gyrA* and *parC* genes were determined by PCR and, if indicated by sequence or restriction analysis [29,30,31,32,33,34,35,36,37].

Characterization of ESBL Producing Isolates

Self-transferability of plasmids was tested by mating experiments using azid-resistant recipient *E. coli* strain J⁵³ as previously described [37]. Further characterization was performed by southern blotting, PCR-based replicon typing and pulsed-field gel electrophoresis (PFGE) using a CHEF DRIII System (BioRad, Munich, Germany) for comparative analysis with clinical isolates [37].

Multilocus Sequence Typing and Phylogenetic Grouping

Multilocus sequence typing (MLST) was carried out for the multiresistant *E. coli* strains according to the scheme developed by Wirth et al. (2006) [17]. Gene amplification and sequencing was done by using primers specified at the *E. coli* MLST web site (<http://mlst.ucc.ie/mlst/mlst/dbs/Ecoli>). Sequences were analyzed by the software package RidomSeqSphere (<http://www.ridom.de/>) and STs were either computed automatically or newly assigned in case novel STs have been identified. *E. coli* phylogenetic groups were determined by Structure analysis based on the concatenated sequences of the seven housekeeping genes (<http://pritch.bsd.uchicago.edu/structure>).

Virulence Gene Typing

Multiresistant *E. coli* isolates were examined for the presence of 59 virulence-associated genes (VAGs) linked with extraintestinal pathogenic and Shiga toxin producing *E. coli* by multiplex and single PCRs as described previously [37]. VAGs determined encode factors within the categories of toxins, adhesins, iron acquisition systems, protectins and others (detailed information is given in Fig. 1).

Chicken Infection Model

Based on phenotypic and genotypic resistance pattern, MLST and virulence gene typing, one exemplary rat isolate (IMT20717; O18:NM:K1; ST95; ST complex 95), resembling a highly virulent ExPEC genotype and serogroup and additionally expressing a CTX-M-9-type beta lactamase, was selected to assess its pathogenic potential *in vivo*.

A chicken infection model [38], which has already been shown to be appropriate for testing non-avian ExPEC strains as well [39] was used. Four groups of six chickens each were infected intratracheally with 10⁹ colony-forming units (CFU) of the test strain and two control strains, one low pathogenic avian fecal strain (IMT12226; O77:H18; ST1165, STC144) as negative control and archetypical *E. coli* strain RS218 (O18:H7:K1; ST95; STC95), isolated from a case of meningitis in a baby, as positive control strain [40]. This non-ESBL producing strain was included in order to directly compare a clinical strain of the same phylogenetic background and identical O- and K-antigens with the multiresistant urban rat isolate.

Results and Discussion

High Isolation Rates of *E. coli* from Rats

E. coli isolates were recovered from 77.0% (n = 67) of the 87 rats (*E. norvegicus*). Isolation from fresh fecal swabs (95%) seemed more preferable than that from swabs taken to the laboratory in conservation media (70%). Overall, the isolation rate is beyond what has been described for other rodent fecal samples so far [41], whereas it is similar to what has been found in a study including wild *R. rattus* in Africa [42]. After picking at least three colonies per animal, a total of 238 *E. coli* isolates were obtained. RAPD-PCR analysis (data not shown) was further used to exclude isolates with identical band profiles, which were only observed in individual

animals but not in-between rats. Finally, 211 non-duplicate strains were included in further experiments and the reduction by a number of only 27 copy strains indicated a quite high diversity amongst rat *E. coli* isolates.

Frequent Occurrence of Multiresistant *E. coli* Isolates in Rats

Several rat *E. coli* isolates showed phenotypic resistance to ampicillin (15%), cephalothin (5%), as well as to fluoroquinolones like enrofloxacin (7%), difloxacin (9%) and marbofloxacin (7%), the two aminoglycosides gentamicin/kanamycin (both 5%), tetracycline (16%), sulfamethoxazole-trimethoprim (9.4%), and chloramphenicol (10%). A total of 55 of all 211 isolates (26%) exhibited resistant phenotypes in the agar dilution test against at least one antimicrobial class. According to MIC data, 32 of these isolates showed resistance to three or more classes of antimicrobials (Table 1). Thus, following the definition given by Schwarz et al. (2010) [43], overall 13.6% (n = 32) of *E. coli* strains isolated from urban rats should be regarded multiresistant. Of these, two (IMT19205 and IMT20717) showed a positive confirmatory test for the production of ESBLs. One of these isolates (IMT19205) belonged to the pandemic clonal group B2-ST131-O25b:H4 and was included in a previous publication [22]. MIC testing of all 32 isolates revealed high rates of resistance to beta-lactams like ampicillin (87.5%), oxacillin (96.9%), cephalothin (31.3%), as well as to fluoroquinolones like enrofloxacin (43.8%), difloxacin (50%) and marbofloxacin (43.8%), the two aminoglycosides gentamicin/kanamycin (both 34.4%), tetracycline (84.4%), sulfamethoxazole-trimethoprim (59.4%), and chloramphenicol (65.6%). The most abundant pattern observed was combined resistance to ampicillin, tetracycline and the fluoroquinolones (Table 1). Screening for antimicrobial resistance determinants nearly always reflected the phenotypic resistance situation. Most or all strains harboured *bla*_{TEM-1-like} (87.5%), *sul1/sul2* (75%) and *strA/B* genes (100%), whereas other non-beta-lactam resistance genes, such as *aadA* (34.4%), *tet(A-D)* (25%), *aac(3)IV* (3.1%), *aac(6')-Ib-cr* (6.3%), and *qnrB1* (3.1%) were present in lower frequencies (Table 1).

In general, data on antimicrobial resistance in *E. coli* from wild rats are rather limited. Literak et al. (2009) identified 2.5% of African *R. rattus* isolates to be ESBL-producers [42]. An additional study reported high rates of multiresistant *E. coli* in rats (*R. norvegicus*) from a port in Greece [6]. Taking into account other synanthropic wildlife species as well, the rates of antimicrobial resistant *E. coli* detected in this study are higher than what has been found in raccoons (16% from urban environments) [44] or small mammals (15% in residential areas) [45]. The higher rates obtained from urban rats could be explained by the assumption that human activities including production of sewage are the most likely common source of *E. coli* transmission to urban wildlife. As only rats populate the sewage system directly, they have direct contact with human feces, whether from private households or clinics and might frequently take up multiresistant strains in this way. The recent finding of comparable antimicrobial resistance patterns in *E. coli* isolates from rats and humans agrees with this [46,47]. Also, compared to rodents from rural areas in Central Europe, the rates of multiresistant *E. coli* from urban rats seem to be higher (13.6% vs. approx. 2%) [22,48]. One logical conclusion could be that rats might serve as surrogate marker for the spread of antimicrobial resistance in urban areas. Above all, however, their potential to disseminate multiresistant microorganisms in highly populated areas should not be obscured, especially since there is almost no doubt about their ability to spread zoonotic pathogens among humans and animals [3,4,5,49].

Table 1. Results of genotypic and phenotypic characterisation of 32 multiresistant rat *E. coli* isolates.

Strain No.	PG	ST	STC	No. VAGs	Phenotypic resistance for antimicrobial substances [no. of classes with phenotypic resistance]	Mutated amino acids encoded in*				Antimicrobial resistance genes/gene variants
						<i>gyrA</i>		<i>parC</i>		
						Ser83	Asp87	Ser80	Glu84	
IMT19258	B2	127	127	31	AMP-OXA-ENR-MAF-DIF-ORB-GEN-KAN-TET-TS [5]	WT	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i>
IMT20717	B2	95	95	31	CEF-AMP-OXA-GEN-KAN-TS [3]					<i>bla</i> _{CTX-M-9} , <i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aac(6')</i> - <i>lb-cr</i> , <i>aacA</i>
IMT19198	B2	1851	none	28	AMP-OXA-TET-TS [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i>
IMT19127	B2	428	none	26	AMP-OXA-CTN-TET-TS [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(B)</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19089	B1	88	23	21	AMP-OXA-CTN-MAF-TET [3]	WT	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19256	ABD	57	350	20	AMP-OXA-ENR-MAF-DIF-ORB-TS [3]	WT	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i> , <i>qnrB1</i>
IMT19129	D	38	38	18	AMP-OXA-CTN-GEN-TET-CMP [4]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19205	B2	131	131	18	CEF-AMP-OXA-ENR-MAF-GEN-TET [4]	Leu	Asn	Ile	Val	<i>bla</i> _{CTX-M-9} , <i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aac(6')</i> - <i>lb-cr</i> , <i>aac(3)IV</i>
IMT19265	Ax81	93	168	17	AMP-OXA-ENR-MAF-DIF-ORB-TS [3]	WT	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>bla</i> _{OXA-11} , <i>sul1</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19282	B2	2381	none	17	AMP-OXA-KAN-TET-CMP [4]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(D)</i> , <i>strA</i> , <i>strB</i>
IMT19110	D	1011	none	16	AMP-OXA-ENR-MAF-DIF-ORB-GEN-TET-TS-CMP [6]	Leu	Asn	Ile	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i>
IMT19264	Ax81	93	168	16	AMP-OXA-TET-TS [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>bla</i> _{OXA-11} , <i>sul1</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19142	ABD	1850	none	14	AMP-OXA-CTN-DIF-TET-CMP [4]	Leu	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>aacA</i>
IMT19176	ABD	641	86	13	AMP-OXA-CTN-ENR-DIF-TET-TS-CMP [5]	Leu	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19179	A	1286	none	13	AMP-OXA-KAN-TET-CMP [4]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i>
IMT19188	B1	1049	none	12	AMP-OXA-CTN-TET-CMP [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19249	ABD	1850	none	12	AMP-OXA-ENR-DIF-TET-CMP [4]	Leu	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(D)</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19173	B1	1049	none	11	AMP-OXA-TET-CMP [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19248	ABD	1850	none	11	AMP-AMC-OXA-DIF-TET-TS-CMP [5]	Leu	Asn	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19262	D	501	none	11	OXA-ENR-MAF-DIF-ORB-TS-CMP [4]	WT	WT	Ile	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>strB</i>
IMT19091	A	10	10	10	AMP-OXA-CTN-ENR-MAF-DIF-ORB-GEN-TET-TS-CMP [6]	Leu	Asn	Ile	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19123	D	70	none	10	OXA-TET-TS [3]					<i>strA</i> , <i>strB</i>
IMT19287	Ax81	1849	none	10	MAF-DIF-GEN-CMP [3]	WT	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(D)</i> , <i>sul1</i> , <i>strA</i>
IMT19121	ABD	453	86	9	AMP-OXA-CTN-TET-CMP [3]					<i>strA</i> , <i>strB</i>
IMT19183	B1	1049	none	9	AMP-OXA-TET-CMP [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19285	D	501	none	9	AMP-OXA-ENR-MAF-DIF-ORB-GEN-KAN-TET-TS-CMP [6]	Leu	Asn	Ile	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i>
IMT19114	B2	1444	none	6	AMP-OXA-TET-CMP [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>strA</i> , <i>strB</i>
IMT19126	ABD	641	86	6	AMP-OXA-TET-CMP [3]					<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(D)</i> , <i>strA</i> , <i>strB</i>
IMT19166	Ax81	224	none	6	AMP-OXA-ENR-MAF-DIF-ORB-GEN-KAN-TET-TS-CMP [6]	Leu	Asn	Ile	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i> , <i>aacA</i>
IMT19257	B1	2380	none	6	OXA-TET-CMP [3]					<i>sul2</i> , <i>strA</i> , <i>strB</i>
IMT19269	Ax81	1849	none	6	AMP-OXA-ENR-MAF-DIF-ORB-TET-TS [4]	Leu	WT	WT	WT	<i>bla</i> _{TEM-1-lik} <i>er</i> , <i>tet(D)</i> , <i>sul1</i> , <i>strA</i> , <i>strB</i>

Table 1. Cont.

Strain No.	PG	ST	STC	No. VAGs	Phenotypic resistance for antimicrobial substances [no. of classes with phenotypic resistance]		Mutated amino acids encoded in*				Antimicrobial resistance genes/gene variants				
					AMP-OXA-ENR-MAF-DIF-ORB-GEN-KAN-TET-TS-CMP [6]	Virulence	Ser83	Asp87	Ser80	Glu84	<i>gyrA</i>	<i>parC</i>	<i>bla</i> _{TEM-1-like}	<i>strA</i>	<i>strB</i>
IMT19158	B1	2976	none	5	AMP-OXA-ENR-MAF-DIF-ORB-GEN-KAN-TET-TS-CMP [6]	Virulence	Ser83	Asp87	Ser80	Glu84	<i>gyrA</i>	<i>parC</i>	<i>bla</i> _{TEM-1-like}	<i>strA</i>	<i>strB</i>

Footnote to Table 1, sorted by the number of virulence associated genes (VAGs; column 5).

AM = antimicrobial, PG = ancestral/phylogenetic group, ST = sequence type, STC = sequence type complex, VAGs = virulence associated genes, AMP = ampicillin, AMC = ampicillin/clavulanic acid, CEF = cefotaxim,

CMP = chloramphenicol, CTN = cefotetan, DIF = difloxacin, ENR = enrofloxacin, GEN = gentamicin, KAN = kanamycin, MAF = marbofloxacin, ORB = orbifloxacin, OXA = oxacillin, TET = tetracycline, TS = sulfamethoxazole/trimethoprim.

*empty fields: not tested (fluoroquinolone sensitive strains).

doi:10.1371/journal.pone.0050331.t001

31), most of these belonging to the B2 group and thus presenting typical ExPEC strains.

Nearly all isolates harboured bacterial adhesin genes encoding Type 1 (*fimC*)- and Curli fimbriae (*csgA*). Also the presence of typical ExPEC-related adhesins, such as the heat-resistant agglutinin [*hxA* (28.1%)], iron-regulated hemagglutinin [*iha* (12.5%)], P-fimbriae [*pap* operon genes (9.4%–12.5%)], S-fimbriae [*sfa/foc* (6.3%)], or a recently described ExPEC adhesin [*ea/I* (18.8%)] hinted towards the affiliation of a number of rat strains to the group of ExPEC strains. Iron acquisition genes, such as *chuA* (43.8%), *fyuA*, *iroN*, *irp2* (all 37.5%), *iucD* (28.1%), *iutA* (40.6%), *sitA* (78.1%), and *sitD*_{episomal} (46.9%), which are known to confer fitness advantage and also invasive properties towards *E. coli* residing in the gut or bladder of their host, under certain circumstances being capable of causing infections at various extraintestinal sites [55,56,57], were also frequently detected. The finding of protectin genes like increased serum resistance gene *iss* (53.1%), and invasion-associated K1-capsule encoding gene *neuC* (9.4%), as well as of plasmid-located transfer [*traT* (56.3%)] and outer membrane genes [*ompT* (43.8%)], all of which are highly associated with the virulence of human and avian ExPEC strains [52,58] substantiates our belief, that rats could frequently be asymptotically colonized by ExPEC-like strains and may thus serve as a permanent source of zoonotic *E. coli*. The pathogenic nature of a number of the strains isolated in the present study is further supported by the detection of toxin genes, such as the cytonecrotizing factor *cnf* (12.5%), secreted autotransporter toxin *sat* (9.4%), vacuolating autotransporter toxin *vat* (9.4%), and haemolysin operon genes *hlyA* and *hlyC* (9.4%), which are particularly characteristic for uropathogenic *E. coli* [56]. Apart from the K1-capsule, which is one of the main features of highly invasive ExPEC strains, exemplified by a subgroup of avian pathogenic *E. coli* (APEC) as well as by *E. coli* strains implicated in new-born meningitis (NMEC), we also found other invasion-related factors among the rat strains, including *ibeA* (9.4%), which has a crucial role in the bacterial translocation of the blood brain barrier epithelium and *in vivo* pathogenicity, as previously shown in a rat meningitis and a chicken infection model [59,60].

Successful with our results, recent publications attributed the successful colonization of the healthy gut of humans, dogs, swine, and poultry also to the presence of ExPEC-typical VAGs [53,61,62,63]. The frequent finding of multiresistant ExPEC-like strains among rat samples, however, contradicts the paradigm about an ultimate loss of bacterial fitness due to the maintenance of antibiotic resistance in combination with high levels of virulence [64]. This combination is considered one of the major drivers for the international spread of ESBL clone O25b:H4-B2-ST131, while there are also studies pointing out that this might be only one side of the coin [21,65]. If virulence would be that decisive for the emergence of antimicrobial resistant and highly virulent ExPEC strains, one would expect other clonal groups, such as the B2-ST95 lineage, which accumulates highly invasive, mostly human and avian strains [17,39,52,53], to acquire a multiresistance phenotype, by that amplifying its threat to human and animal health. Though, as discussed earlier, so far only a marginal proportion (4%) of all ST-complex 95 strains deposited on the web-hosted database (<http://mlst.ucc.ie/mlst/mlst/dbs/Ecoli/>) or reported in several publications harbors ESBL genes or simply a multiresistant phenotype [21]. The more intriguing it was that we identified an ST95 ESBL-producing strain (IMT20717; CTX-M-9) among the rat isolates, which remarkably showed multiresistance, frequent possession of virulence genes (n = 31) in a B2 phylogenetic background, and a serogroup (O18:NM:K1) typical of highly invasive ExPEC strains (Fig. 1; Table 1). In that way it




Figure 2. Dendrogram of ST95-ESBL rat strain IMT20717 with *E. coli* ST95-K1 strains. The clonal relationship shown is based on *Xba*I-generated PFGE profiles. NM=non motile (H antigen negative or not expressed); NBM=newborn meningitis; UTI=urinary tract infection, optimization 1.0%, position tolerance 1.5%. doi:10.1371/journal.pone.0050331.g002




Figure 3. Results of the ST95-ST95 ESBL rat strain IMT20717 in the chicken infection model. Ability of B2-ST95-O18:NM:K1-CTX-M-9 urban rat strain IMT20717 to colonize the lungs, disseminate into internal organs and penetrate the blood brain barrier 24 h post intra-tracheal infection (10^9 CFU) of a group of six 5-weeks old SPF White Leghorn chickens. Non-ESBL-producing NMEC strain RS218 (B2-ST95-O18:H7:K1) and avian fecal strain IMT12226 (ST1165-O77:H18), known invasive and low pathogenic strains, were used as controls. doi:10.1371/journal.pone.0050331.g003

very much resembles isolates causing urosepsis and new born meningitis in humans, and septicemia in chickens. Due to its observed lifestyle in the rat, namely asymptotically colonizing the gut, it was reasonable to deduce the strains extraintestinal pathogenicity not simply from its phylogenetic background and the possession of several VAGs, but also experimentally in an *in vivo* model. We made use of chicken experiments as this has been shown a proper model for determining the pathogenicity of ExPEC strains, in particular of ST95 strains, which are highly linked to chickens as one of their natural hosts [39].

Paradigmatic Combination of Multiresistance and Extraintestinal Pathogenicity in Urban Rat ST95-CTX-M-9-producing Strain IMT20717

IMT20717 displayed a positive confirmatory test for the production of ESBL. Apart from a *bla*_{CTX-M-9} gene, this strain also harboured resistance genes *bla*_{TEM-1}, *sul2*, *strA*, *strB*, *aac(6′)-Ib-cr* and *aadA*. All these genes, except for *sul2*, were located on a self-transferable, approximately 50 kb plasmid of the N/FIC replicon type.

Serotyping characterized IMT20717 as O18:NM:K1. Thereby the strain expressed a combination of an O-antigen and a capsule type which is highly linked with a clonal group of *E. coli* strains frequently involved in invasive infections in humans [19,39,58,59,66]. Particular attention has been drawn to *E. coli* O18:K1 NMEC strains causing meningitis in babies shortly after delivery. In addition, this serogroup is also frequent among avian pathogenic *E. coli* (APEC). Here, it causes often fatal septicemia and is responsible for great losses in poultry breeding [39,52,66]. Macrorestriction analysis and subsequent PFGE revealed a high genetic similarity (Dice similarity $\geq 82.2\%$) between the rat B2-ST95-O18:NM:K1-CTX-M-9 isolate and clinical ST95 strains of different ExPEC pathovars, and an additional fecal strain from the gut of a healthy human, all affiliated to this globally distributed lineage (Fig. 2). This similarity strongly resembles what is already well known, in that, the healthy human gut serves as a reservoir for these pathovars [61]. The detection of a pathogenic strain linked to human clinical environments points towards a possible transmission pathway through clinic waste into the urban sewage system.

In the *in vivo* infection model IMT20717 revealed a lower bacterial recovery rate from chicken organs than the clinical NMEC type strain RS218 (Fig. 3), which was included for comparative purposes. Nevertheless the strain could be isolated from all internal organs in significantly higher numbers than the low pathogenic avian control strain IMT12226. Particularly its re-isolation from the brain is of high indicative value for its invasive potential (Fig. 3) as it suggests that the strain is able to penetrate the blood brain barrier. In view of this, the finding of an ST95 strain in the gut of a rat could add rats to the list of potential hosts for highly extraintestinal pathogenic *E. coli* as well. What is even

more disturbing is the fact that this strain carried an ESBL-encoding plasmid. Such a combination, namely ESBL-ST95-*E. coli*, has only rarely been observed in human clinical samples so far [21] and its detection in wild rat is the first description of such a superbug in an animal.

Conclusions

The urban rats examined in this study frequently carried multiresistant *E. coli* strains showing high levels of resistance to critically important antimicrobials like fluoroquinolones and β -lactams. As the WHO classified urban rats as a significant public health threat [2] the data reported here might have yet unpredictable consequences in the future. In particular, the finding of an ESBL-producing ExPEC strain belonging to one of the most virulent ExPEC lineages (ST95) might signify a new development in the field of antimicrobial resistance, in that ESBL plasmids step by step could find their way into highly virulent *E. coli* populations. There is still no final clue for the recent dominance of the pandemic B2-ST131-O25b:H4-CTX-M-15 clonal group. Several non-resistance-related attributes like bacterial fitness, virulence patterns or insertional modifications in fimbrial genes have been discussed as putative causes [65,67,68]. Taking into account the virulence potential of ST95, which is believed to be comparably high to that of ST131, it remains unclear why ST95 is far from being as broadly distributed as ESBL-producing ST131. However, the urban rat-derived B2-ST95-O18:NM:K1-CTX-M-9 strain possesses a number of genetic markers whose products confer adhesive, toxic and invasive properties and thus meets all requirements for a successful commensal and extraintestinal pathogenic life style. Future monitoring of clinical and environmental ESBL *E. coli* isolates should therefore clarify whether the detection of this ST95-ESBL strain from a rat simply presents an accidental finding of a minor important ESBL clone in a single animal, or whether it points towards a successful spread of ST95-ESBL outside the clinics as well. In any case, this strain hallmarks the main finding of this study: the mere occurrence of *E. coli* strains in urban rats that are multiresistant & virulent is an alarming observation, as infections with such strains could lead to severe clinical outcomes, leaving only limited treatment options.

Acknowledgments

We would like to thank Matthias Stange and Mario Heising for providing their excellent assistance in rat sampling, Ines Diehl for her excellent technical assistance, and Yvonne Pfeifer for providing *E. coli* control strains.

Author Contributions

Conceived and designed the experiments: SG LHW CE. Performed the experiments: SG AB AF TS RGU CE. Analyzed the data: SG LHW TS CE. Wrote the paper: SG LHW CE.

References

1. Nkogwe C, Raletobana J, Stewart-Johnson A, Suepaul S, Adesiyun A (2011) Frequency of Detection of *Escherichia coli*, *Salmonella* spp., and *Campylobacter* spp. in the Faeces of Wild Rats (*Rattus* spp.) in Trinidad and Tobago. *Vet Med Int*: doi:10.4061/2011/686923.
2. Bonnefoy X, Kampen H, Sweeney K (2008) Public Health Significance of Urban Pests. WHO Regional Office for Europe.
3. Meerburg BG, Singleton GR, Kijlstra A (2009) Rodent-borne diseases and their risks for public health. *Crit Rev Microbiol* 35: 221–270.
4. Runge M, von Keyserlingk M, Braune S, Becker D, Plenge-Bonig A, et al. (2012) Distribution of rodenticide resistance and zoonotic pathogens in Norway rats in Lower Saxony and Hamburg, Germany. *Pest Manag Sci*.
5. Cizek A, Alexa P, Literak I, Hamrik J, Novak P, et al. (1999) Shiga toxin-producing *Escherichia coli* O157 in feedlot cattle and Norwegian rats from a large-scale farm. *Lett Appl Microbiol* 28: 435–439.
6. Burriel AR, Kritas SK, Kontos V (2008) Some microbiological aspects of rats captured alive at the port city of Piraeus, Greece. *Int J Environ Health Res* 18: 159–164.
7. Gardner-Santana LC, Norris DE, Fornadel CM, Hinson ER, Klein SL, et al. (2009) Commensal ecology, urban landscapes, and their influence on the genetic characteristics of city-dwelling Norway rats (*Rattus norvegicus*). *Molecular Ecology* 18: 2766–2778.
8. Orgain H, Schein MW (1953) A Preliminary Analysis of the Physical Environment of the Norway Rat. *Ecology* 34: 467–473.

9. Emlen JT Jr, Davis DE (1948) Determination of reproductive rates in rat populations by examination of carcasses. *Physiol Zool* 21: 59–65.
10. Battersby SA, Parsons R, Webster JP (2002) Urban rat infestations and the risk to public health. *J Environ Health Res* 1: 4.
11. Channon D, Cole M, Cole L (2000) A long-term study of *Rattus norvegicus* in the London Borough of Enfield using baiting returns as an indicator of sewer population levels. *Epidemiol Infect* 125: 441–445.
12. Davis DE, Hall O (1951) The seasonal reproductive condition of female Norway (brown) rats in Baltimore, Maryland. *Physiological Zoology* 24.
13. Taylor KD, Quyy RJ (1978) Long distance movements of a common rat (*Rattus norvegicus*) revealed by radio-tracking. *Mammalia* 42: 47–53.
14. Glass GE, Childs JE, Korch GW, LeDuc JW (1989) Comparative ecology and social interactions of Norway rat (*Rattus norvegicus*) populations in Baltimore. Occasional Papers of the Museum of Natural History, University of Kansas, Lawrence, Kansas 130.
15. Webster JP, Macdonald DW (1995) Parasites of Wild Brown-Rats (*Rattus Norvegicus*) on Uk Farms. *Parasitology* 111: 247–255.
16. Meyer AN, Shankster A, Langton SD, Jukes G (1995) National Commensal Rodent Survey 1993. *Environmental Health* 103: 127–135.
17. Wirth T, Falush D, Lan R, Colles F, Mensa P, et al. (2006) Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 60: 1136–1151.
18. Nielsen EM, Skov MN, Madsen JJ, Lodal J, Jespersen JB, et al. (2004) Verocytotoxin-producing *Escherichia coli* in wild birds and rodents in close proximity to farms. *Appl Environ Microbiol* 70: 6944–6947.
19. Dubois D, Prasadarao NV, Mittal R, Bret L, Roujou-Gris M, et al. (2009) CTX-M beta-lactamase production and virulence of *Escherichia coli* K1. *Emerg Infect Dis* 15: 1988–1990.
20. Pitout JD (2012) Extraintestinal Pathogenic *Escherichia coli*: A Combination of Virulence with Antibiotic Resistance. *Front Microbiol* 3: 9.
21. Ewers C, Bethe A, Semmler T, Guenther S, Wieler LH (2012) Extended-spectrum beta-lactamase-producing and AmpC-producing *Escherichia coli* from livestock and companion animals, and their putative impact on public health: a global perspective. *Clin Microbiol Infect*.
22. Guenther S, Grobbel M, Beutlich J, Guerra B, Ulrich RG, et al. (2010) Detection of pandemic B2-O25-ST131 *Escherichia coli* harbouring the CTX-M-9 extended-spectrum beta-lactamase type in a feral urban brown rat (*Rattus norvegicus*). *J Antimicrob Chemother* 65: 582–584.
23. Platell JL, Johnson JR, Cobbold RN, Trott DJ (2011) Multidrug-resistant extraintestinal pathogenic *Escherichia coli* of sequence type ST131 in animals and foods. *Vet Microbiol* 153: 99–108.
24. Rogers BA, Sidjabat HE, Paterson DL (2011) *Escherichia coli* O25b-ST131: a pandemic, multiresistant, community-associated strain. *J Antimicrob Chemother* 66: 1–14.
25. Winkle S (1979) Mikrobiologische und serologische Diagnostik. Jena: VEB Gustav Fischer Verlag.
26. Schierack P, Römer A, Jores J, Kaspar H, Guenther S, et al. (2008) Isolation and characterization of intestinal *E. coli* from wild boars in Germany. *Appl Environ Microbiol* 75: 695–702.
27. Guenther S, Grobbel M, Lubke-Becker A, Goedecke A, Friedrich ND, et al. (2010) Antimicrobial resistance profiles of *Escherichia coli* from common European wild bird species. *Vet Microbiol* 144: 219–225.
28. CLSI (2008) Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; approved standard - Third Edition. Wayne, PA, U.S.A.: CLSI.
29. Rodriguez I, Barownick W, Helmuth R, Mendoza MC, Rodicio MR, et al. (2009) Extended-spectrum beta-lactamases and AmpC beta-lactamases in cefitour-resistant *Salmonella enterica* isolates from food and livestock obtained in Germany during 2003–07. *J Antimicrob Chemother* 64: 301–309.
30. Bertrand S, Weill FX, Cloeckaert A, Vrints M, Mairiaux E, et al. (2006) Clonal emergence of extended-spectrum beta-lactamase (CTX-M-2)-producing *Salmonella enterica* serovar Virchow isolates with reduced susceptibilities to ciprofloxacin among poultry and humans in Belgium and France (2000 to 2003). *J Clin Microbiol* 44: 2897–2903.
31. Park YJ, Yu JK, Lee S, Oh EJ, Woo GJ (2007) Prevalence and diversity of *qnr* alleles in AmpC-producing *Enterobacter cloacae*, *Enterobacter aerogenes*, *Citrobacter freundii* and *Serratia marcescens*: a multicentre study from Korea. *J Antimicrob Chemother* 60: 868–871.
32. Robicsek A, Strahilevitz J, Jacoby GA, Macielag M, Abbanat D, et al. (2006) Fluoroquinolone-modifying enzyme: a new adaptation of a common aminoglycoside acetyltransferase. *Nat Med* 12: 83–88.
33. Grimm V, Ezaki S, Susa M, Knabbe C, Schmid RD, et al. (2004) Use of DNA microarrays for rapid genotyping of TEM beta-lactamases that confer resistance. *J Clin Microbiol* 42: 3766–3774.
34. Pomba C, da Fonseca JD, Baptista BC, Correia JD, Martinez-Martinez L (2009) Detection of the pandemic O25-ST131 human virulent *Escherichia coli* CTX-M-15-producing clone harboring the *qnrB2* and *aac(6′)-Ib-cr* genes in a dog. *Antimicrob Agents Chemother* 53: 327–328.
35. Ewers C, Bethe A, Wieler LH, Guenther S, Stamm I, et al. (2011) Companion animals: a relevant source of extended-spectrum beta-lactamase-producing fluoroquinolone-resistant *Citrobacter freundii*. *Int J Antimicrob Agents* 37: 86–87.
36. Costa D, Poeta P, Saenz Y, Vinue L, Coelho AC, et al. (2008) Mechanisms of antibiotic resistance in *Escherichia coli* isolates recovered from wild animals. *Microb Drug Resist* 14: 71–77.
37. Ewers C, Grobbel M, Stamm I, Kopp PA, Diehl I, et al. (2010) Emergence of human pandemic O25:H4-ST131 CTX-M-15 extended-spectrum beta-lactamase-producing *Escherichia coli* among companion animals. *J Antimicrob Chemother* 65: 651–660.
38. Antao EM, Glodde S, Li G, Sharifi R, Homeier T, et al. (2008) The chicken as a natural model for extraintestinal infections caused by avian pathogenic *Escherichia coli* (APEC). *Microb Pathog* 45: 361–369.
39. Moulin-Schouleur M, Reperant M, Laurent S, Bree A, Mignon-Grasteau S, et al. (2007) Extraintestinal pathogenic *Escherichia coli* strains of avian and human origin: link between phylogenetic relationships and common virulence patterns. *J Clin Microbiol* 45: 3366–3376.
40. Silver RP, Aaronson W, Sutton A, Schneerson R (1980) Comparative analysis of plasmids and some metabolic characteristics of *Escherichia coli* K1 diseased and healthy individuals. *Infect Immun* 29: 200.
41. Swiecicka I, Buczek J, Iwaniuk A (2003) Analysis of genetic relationships and antimicrobial susceptibility of *Escherichia coli* isolated from *Clethrionomys glareolus*. *J Gen Appl Microbiol* 49: 315–320.
42. Literak I, Dolejska M, Cizek A, Djigo CAT, Konecny A, et al. (2009) Reservoirs of antibiotic-resistant *Enterobacteriaceae* among animals sympatric to humans in Senegal: extended-spectrum beta-lactamases in bacteria in a black rat (*Rattus rattus*) Afri J Microbiol Res 3: 751–754.
43. Schwarz S, Silley P, Simjee S, Woodford N, van Duijkeren E, et al. (2010) Editorial: assessing the antimicrobial susceptibility of bacteria obtained from animals. *J Antimicrob Chemother* 65: 601–604.
44. Jardine CM, Janecko N, Allan M, Boerlin P, Chalmers G, et al. (2012) Antimicrobial resistance in *Escherichia coli* isolates from raccoons (*Procyon lotor*) in Southern Ontario, Canada. *Appl Environ Microbiol* 78: 3873–3879.
45. Allen SE, Boerlin P, Janecko N, Lumsden JS, Barker IK, et al. (2011) Antimicrobial resistance in generic *Escherichia coli* isolates from wild small mammals living in swine farm, residential, landfill, and natural environments in southern Ontario, Canada. *Appl Environ Microbiol* 77: 882–888.
46. Dhanji H, Murphy NM, Akhigbe C, Doumth M, Hope R, et al. (2011) Isolation of fluoroquinolone-resistant O25b:H4-ST131 *Escherichia coli* with CTX-M-14 extended-spectrum beta-lactamase from UK river water. *J Antimicrob Chemother* 66: 512–516.
47. Guenther S, Ewers C, Wieler LH (2011) Extended-spectrum beta-lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front Microbiol* 2.
48. Guenther S, Grobbel M, Heidemanns K, Schlegel M, Ulrich RG, et al. (2010) First insights into antimicrobial resistance among faecal *Escherichia coli* isolates from small wild mammals in rural areas. *Sci Total Environ* 408: 3519–3522.
49. Nijsten R, London N, van den Bogaard A, Stobberingh E (1995) In-vivo transfer of resistance plasmids in rat, human or pig-derived intestinal flora using a rat model. *J Antimicrob Chemother* 36: 975–985.
50. Bukh AS, Schonheyder HC, Emmersen JMG, Sogaard M, Bastholm S, et al. (2009) *Escherichia coli* phylogenetic groups are associated with site of infection and level of antibiotic resistance in community-acquired bacteraemia: a 10 year population-based study in Denmark. *J Antimicrob Chemother* 64: 163–168.
51. Johnson JR, Johnston B, Clabots C, Kuskowski MA, Pendyala S, et al. (2010) *Escherichia coli* sequence type ST131 as an emerging fluoroquinolone-resistant uropathogen among renal transplant recipients. *Antimicrob Agents Chemother* 54: 546–550.
52. Johnson TJ, Wannemuehler Y, Johnson SJ, Stell AL, Doetkott C, et al. (2008) Comparison of extraintestinal pathogenic *Escherichia coli* strains from human and avian sources reveals a mixed subset representing potential zoonotic pathogens. *Appl Environ Microbiol* 74: 7043–7050.
53. Ewers C, Antao EM, Diehl I, Philipp HC, Wieler LH (2009) Intestine and environment of the chicken as reservoirs for extraintestinal pathogenic *Escherichia coli* strains with zoonotic potential. *Appl Environ Microbiol* 75: 184–192.
54. Sargeant JM, Sanderson MW, Griffin DD, Smith RA (2004) Factors associated with the presence of *Escherichia coli* O157 in feedlot-cattle water and feed in the Midwestern USA. *Prev Vet Med* 66: 207–237.
55. Johnson JR, Russo TA (2005) Molecular epidemiology of extraintestinal pathogenic (uropathogenic) *Escherichia coli*. *Int J Med Microbiol* 295: 383–404.
56. Wiles TJ, Kulesus RR, Mulvey MA (2008) Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol* 85: 11–19.
57. Feldmann F, Sorsa IJ, Hildinger K, Schubert S (2007) The salmochelin siderophore receptor *IronN* contributes to invasion of urothelial cells by extraintestinal pathogenic *Escherichia coli* in vitro. *Infect Immun* 75.
58. Ewers C, Li G, Wilking H, Kiessling S, Alt K, et al. (2007) Avian pathogenic, uropathogenic, and newborn meningitis-causing *Escherichia coli*: how closely related are they? *Int J Med Microbiol* 297: 163–176.
59. Kim BY, Kang J, Kim KS (2005) Invasion processes of pathogenic *Escherichia coli*. *Int J Med Microbiol* 295: 463–470.
60. Germon P, Chen YH, He L, Blanco JE, Bree A, et al. (2005) *ibeA*, a virulence factor of avian pathogenic *Escherichia coli*. *Microbiology* 151: 1179–1186.
61. Nowrouzian FL, Adlerberth I, Wold AE (2006) Enhanced persistence in the colonic microbiota of *Escherichia coli* strains belonging to phylogenetic group B2: role of virulence factors and adherence to colonic cells. *Microbes Infect* 8: 834–840.
62. Johnson JR, Stell AL, Delavari P (2001) Canine feces as a reservoir of extraintestinal pathogenic *Escherichia coli*. *Infect Immun* 69: 1306–13114.

63. Schierack P, Walk N, Ewers C, Wilking H, Steinruck H, et al. (2008) ExPEC-typical virulence-associated genes correlate with successful colonization by intestinal *E. coli* in a small piglet group. *Environ Microbiol* 10: 1742–1751.
64. Johnson JR, Johnston B, Clabots C, Kuskowski MA, Castanheira M (2010) *Escherichia coli* sequence type ST131 as the major cause of serious multidrug-resistant *E. coli* infections in the United States. *Clin Infect Dis* 51: 286–294.
65. Croxall G, Hale J, Weston V, Manning G, Cheetham P, et al. (2011) Molecular epidemiology of extraintestinal pathogenic *Escherichia coli* isolates from a regional cohort of elderly patients highlights the prevalence of ST131 strains with increased antimicrobial resistance in both community and hospital care settings. *J Antimicrob Chemother* 66: 2501–2508.
66. Achtman M, Pluschke G (1986) Clonal analysis of descent and virulence among selected *Escherichia coli*. *Annu Rev Microbiol* 40: 185–210.
67. Pitout JDD (2010) Infections with extended-spectrum beta-lactamase-producing *Enterobacteriaceae*: changing epidemiology and drug treatment choices. *Drugs* 70: 313–333.
68. Totsika M, Beatson SA, Sarkar S, Phan MD, Petty NK, et al. (2011) Insights into a multidrug resistant *Escherichia coli* pathogen of the globally disseminated ST131 lineage: genome analysis and virulence mechanisms. *PLoS One* 6: e26578.