

Journal of Health Monitoring · 2020 5(S7)

DOI 10.25646/7134

Robert Koch Institute, Berlin

Benjamin Wachtler<sup>1</sup>, Niels Michalski<sup>1</sup>, Enno Nowossadeck<sup>1</sup>, Michaela Diercke<sup>2</sup>, Morten Wahrendorf<sup>3</sup>, Claudia Santos-Hövenner<sup>1</sup>, Thomas Lampert<sup>1</sup>, Jens Hoebel<sup>1</sup>

<sup>1</sup> Robert Koch Institute, Berlin

Department of Epidemiology and Health Monitoring

<sup>2</sup> Robert Koch Institute, Berlin

Department of Infectious Disease Epidemiology

<sup>3</sup> University of Düsseldorf, Medical Faculty, Institute of Medical Sociology, Centre for Health and Society

Please cite this publication as

Wachtler B, Michalski N, Nowossadeck E, Diercke M, Wahrendorf M et al. (2020) Socioeconomic inequalities and COVID-19 – A review of the current international literature. Additional material: Overview of the included publications.

Journal of Health Monitoring 5(S7): e1-e12. DOI 10.25646/7134

**Socioeconomic inequalities and COVID-19 – A review of the current international literature. Additional material: Overview of the included publications\***

Source: Own table

| Peer-reviewed articles | # | Author              | Title                                                                                                                                                                                           | Country | Date published | Study design         | Study population                                                                                 | Study period        | Measures of Socioeconomic Status                                                                                  | Outcome measures           | Results                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|------------------------|---|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|----------------|----------------------|--------------------------------------------------------------------------------------------------|---------------------|-------------------------------------------------------------------------------------------------------------------|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                        | 1 | Azar et al. [41]    | Disparities In Outcomes Among COVID-19 Patients In A Large Health Care System In California                                                                                                     | USA     | 21.05.2020     | Retrospective cohort | 14,036 tested, 1,052 confirmed cases of one integrated health care system in Northern California | 01.01. - 08.04.2020 | Ethnicity/race, income, insurance plan                                                                            | Incidence, hospitalisation | Individuals with Medicaid or who were self-pay or had no reported insurance had twice the odds of being admitted, compared to those with commercial insurance (OR=2.1 for both; $p<0,05$ ). COVID-19 positive patients residing in ZIP codes within the top two quartiles of income (quartiles 3 and 4) were less likely to be admitted to the hospital than those residing in the bottom-quartile ZIP code (OR=0.24 and 0.55 for the top two quartiles).                                                           |
|                        | 2 | Kim & Bostwick [65] | Social Vulnerability and Racial Inequality in COVID-19 Deaths in Chicago                                                                                                                        | USA     | 21.05.2020     | Ecological study     | 269 deaths in Chicago                                                                            |                     | Ethnicity/race, Social Vulnerability Index (SVI)                                                                  | Mortality                  | Community areas with higher levels of SVI and risk factor score had a significantly higher COVID death rate. For the most part, the effect of social vulnerability on the death rate was mediated through health risk factors.                                                                                                                                                                                                                                                                                      |
|                        | 3 | Lassale et al. [68] | Ethnic disparities in hospitalisation for COVID-19 in England: The role of socioeconomic factors, mental health, and inflammatory and proinflammatory factors in a community-based cohort study | England | 01.06.2020     | Retrospective cohort | 340,966 participants, 640 cases                                                                  | 16.03. - 26.04.2020 | Self-reported ethnicity, Townsend index of area deprivation, household income, educational attainment, occupation | Hospitalisation            | After adjusting for age and sex, compared to White participants, being from a Black ethnic background was associated with over a four-fold risk of hospitalisation for COVID-19 (odds ratio; 95% confidence interval: 4.32; 3.00–6.23), while a doubling was apparent in Asian (2.12; 1.37, 3.28) and other ethnic groups (1.84; 1.13, 2.99). The greatest attenuations were observed when socioeconomic factors were added to the multivariable model: 24.5% for Blacks, 31.9.3% for Asians, and 30.0% for others. |

\* References can be found in the article [Wachtler B, Michalski N, Nowossadeck E, Diercke M, Wahrendorf M et al. \(2020\) Socioeconomic inequalities and COVID-19 – A review of the current international literature. Journal of Health Monitoring 5\(S7\): 3–17](#) of the Journal of Health Monitoring.

|   |                           |                                                                                                         |         |            |                      |                                                                                   |                     |                                                                                                        |                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|---|---------------------------|---------------------------------------------------------------------------------------------------------|---------|------------|----------------------|-----------------------------------------------------------------------------------|---------------------|--------------------------------------------------------------------------------------------------------|------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 4 | Millett et al. [18]       | Assessing Differential Impacts of COVID-19 on Black Communities                                         | USA     | 14.05.2020 | Ecological study     | US population                                                                     | Until 13.04.2020    | Ethnicity/race, insurance, household crowding, unemployment                                            | Incidence, mortality               | Higher rates of COVID-19 cases were independently associated with greater proportions of uninsured residents (RR 1.16, 95% CI 1.07-1.126), higher percentages of residents in crowded living conditions (RR 1.05, 95% CI 1.01-1.10), and more days since the first case (RR 3.1, 95% CI 2.9-3.3). 280,112 cases of excess COVID-19 diagnoses were associated with occupancy of greater than one person per room, and 126,985 excess diagnoses of COVID-19 were associated with lack of health insurance. The population attributable fraction (PAF) for lack of health insurance was 3.3% for counties with <13% black residents and 4.2% for counties with >13% black residents. Higher county-level unemployment was associated with fewer COVID-19 diagnoses. |
| 5 | Mollalo et al. [42]       | GIS-based spatial modeling of COVID-19 incidence rate in the continental United States                  | USA     | 22.04.2020 | Ecological study     | US population                                                                     | 22.01. - 09.04.2020 | Median household income, income inequalities, percentage uninsured, unemployment rate, food insecurity | Incidence                          | Income inequality, median household income, the percentage of nurse practitioners, and the percentage of the black female population (to the total female population) at the county-level were positively associated with the COVID-19 incidence rate on county level.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| 6 | Niedzwiedz et al. [67]    | Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank | England | 29.05.2020 | Retrospective cohort | 392,116 participants in England, 2,658 tested, 948 cases, 726 in hospital         | 16.03 - 03.05.2020  | Ethnicity/race, occupation, Townsend Index of area deprivation, educational level, employment status   |                                    | Living in a disadvantaged area (according to the Townsend deprivation score) was associated with a higher risk of confirmed infection, particularly for the most disadvantaged quartile (RR 2.19 (95% CI 1.80–2.66)). Differences in ethnicity and country of birth, social factors, baseline health and behavioural risk factors all moderately attenuated the association in the most disadvantaged quartile.                                                                                                                                                                                                                                                                                                                                                  |
| 7 | Okoh et al. [83]          | Coronavirus disease 19 in minority populations of Newark, New Jersey                                    | USA     | 10.06.2020 | Retrospective cohort | African Americans and Hispanics admitted to one hospital in Newark, n = 416 cases | 10.03. - 10.04.2020 | Population densities, housing units and median income on zip code level                                | Hospitalisation, mortality         | Minority patients admitted for COVID-19 were in the highest quartile for population's density, number of housing units and disproportionately fell into the lowest median income quartile.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| 8 | Price-Haywood et al. [40] | Hospitalization and Mortality among Black Patients and White Patients with Covid-19                     | USA     | 27.05.2020 | Retrospective cohort | 3,481 cases                                                                       | 01.03. - 11.04.2020 | Ethnicity/race, insurance plan, zip code level income                                                  | Length of hospital stay, mortality | Increasing age, a higher score on the Charlson Comorbidity Index, public insurance (Medicare or Medicaid), residence in a low-income area, and obesity were associated with an increased odds of admission. However, black race was not independently associated with risk of in-hospital death (hazard ratio, 0.89; 95% CI, 0.68 to 1.17).                                                                                                                                                                                                                                                                                                                                                                                                                      |

|  |    |                     |                                                                                             |     |            |                  |                             |                     |                                  |                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|--|----|---------------------|---------------------------------------------------------------------------------------------|-----|------------|------------------|-----------------------------|---------------------|----------------------------------|---------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | 9  | Ramirez & Lee [53]  | COVID-19 Emergence and Social and Health Determinants in Colorado: A Rapid Spatial Analysis | USA | 29.05.2020 | Ecological study | Population of Colorado      | 14.03. - 08.04.2020 | Social vulnerability index (SVI) | Mortality                             | Higher COVID-19-related death rates were population density and asthma hospitalization, suggestive of urban areas, and poverty and unemployment, suggestive of rural areas. Furthermore, a spatial overlap of high rates of chronic diseases with high rates of COVID-19 may suggest a broader syndemic health burden, where comorbidities intersect with inequality of social determinants of health.                                                                                                                                                                                                                                                                                                               |
|  | 10 | Wadhera et al. [54] | Variation in COVID-19 Hospitalizations and Deaths Across New York City Boroughs             | USA | 29.04.2020 | Ecological study | Population of New York City | Until 25.04.2020    | Median income, Education         | Incidence, hospitalisation, mortality | The number of patients with COVID-19 who were hospitalized per 100,000 population was highest in the Bronx (634) and lowest in Manhattan (331). The number of deaths related to COVID-19 per 100,000 population was also highest in the Bronx (224) and lowest in Manhattan (122). Household median income was lowest in the Bronx (\$38,467) as was the proportion of persons with a bachelor's degree or higher (20.7%). There were 48 short-term acute care hospitals. The number of hospitals per borough ranged from 2 in Staten Island to 16 in Manhattan. The number of hospital beds per 100,000 population was lowest in Queens (144 beds) and highest in the Bronx (336 beds) and in Manhattan (534 beds). |

|                                |    |                   |                                                                                                  |     |            |                  |                                                                             |                     |                                                                                       |                      |                                                                                                                                                                                                                                                                                                                                                                                             |
|--------------------------------|----|-------------------|--------------------------------------------------------------------------------------------------|-----|------------|------------------|-----------------------------------------------------------------------------|---------------------|---------------------------------------------------------------------------------------|----------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Not peer-reviewed publications | 11 | Abedi et al. [43] | Racial, Economic and Health Inequality and COVID-19 Infection in the United States               | USA | 01.05.2020 | Ecological study | Population of 369 counties (total population: 102,178,117 [median: 73,447]) | 05.04. - 16.04.2020 | Ethnicity/race, poverty level, median income, education, rate of uninsured population | Incidence, mortality | Counties with more diverse demographics, higher population, education, income levels, and lower disability rates were at a higher risk of COVID-19 infection. Counties with higher disability and poverty rates had a higher death rate. African Americans were more vulnerable to COVID-19 than other ethnic groups (1,981 African American infected cases versus 658 Whites per million). |
|                                | 12 | Ahmad et al. [81] | Association of Poor Housing Conditions with COVID-19 Incidence and Mortality Across US Counties. | USA | 30.05.2020 | Ecological study | 3,141 US counties                                                           |                     | Poor housing conditions, median household income, education, ethnicity                | Incidence            | In the fully adjusted models, with each 5% increase in percent households with poor housing conditions, there was a 50% higher risk of COVID-19 incidence (IRR 1.50, 95% CI: 1.38 – 1.62) and a 42% higher risk of COVID-19 mortality (MRR 1.42, 95% CI: 1.25 – 1.61). Results remained similar using earlier timepoints (3/31/2020 and 4/10/2020).                                         |

|    |                        |                                                                                                                                                                |         |            |                      |                                                                    |                     |                                                                               |                                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|----|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------|----------------------|--------------------------------------------------------------------|---------------------|-------------------------------------------------------------------------------|----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 13 | Apea et al. [69]       | Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: an observational cohort study                                          | England | 12.06.2020 | Retrospective cohort | 1,737 patients admitted to five acute NHS Hospitals in East London | 01.03. - 13.05.2020 | Ethnicity, Index of Multiple Deprivation                                      | Mortality, admission to intensive care | The majority of patients were classified as being in the two most deprived socio-economic quintiles in England.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| 14 | Buja et al. [62]       | Demographic and Socio-Economic Factors, and Healthcare Resource Indicators Associated with the Rapid Spread of COVID-19 in Northern Italy: An Ecological Study | Italy   | 29.04.2020 | Ecological study     | Population of 36 provinces in Northern Italy                       | 24.02. - 30.03.2020 | Regional employment rates                                                     | Incidence                              | A significant positive correlation was found with employment rates, public transportation rates, in-house density, population density, and the proportions of private acute and long-term care beds in clinics and nursing homes.                                                                                                                                                                                                                                                                                                                                                                                                                     |
| 15 | Chow et al. [44]       | The disproportionate rise in COVID-19 cases among Hispanic/Latinx in disadvantaged communities of Orange County, California: A socioeconomic case-series       | USA     | 07.05.2020 | Ecological study     | 154 cases                                                          | 12.03. - 22.04.2020 | Ethnicity/race, regional household income, education, health insurance status | Incidence                              | Hispanic/Latinx patients residing in census tracts below the median income demonstrated exponential growth (rate = 55.9%, R <sup>2</sup> = 0.9742) during the study period. In addition, there was a significant difference for both race-ethnic (p < 0.001) and income bracket (p = 0.001) distributions prior to and after California's shelter-in-place. In addition, the percentage of individuals residing in neighborhoods with denser households (p = 0.046), lower levels of college graduation (p < 0.001), health insurance coverage (p = 0.01), and ability to work from home (p < 0.001) significantly increased over the same timeframe. |
| 16 | Cyrus et al. [55]      | The impact of COVID-19 on African American communities in the United States                                                                                    | USA     | 19.05.2020 | Ecological study     | 152 counties                                                       | 22.01. - 12.04.2020 | Poverty level                                                                 | Incidence, mortality                   | COVID-19 prevalence increased 5% for every percentage increase in county African American density, and the death rate increased 2 per hundred thousand for every percentage increase in county African American density.                                                                                                                                                                                                                                                                                                                                                                                                                              |
| 17 | Federgruen & Naha [56] | Variation in Covid-19 Cases Across New York City                                                                                                               | USA     | 29.05.2020 | Ecological study     | Population of New York City                                        |                     | Poverty Level                                                                 | Incidence                              | An increase of the average household size by one member accounts, in our final model specification, for at least 876 cases, a full 23% of the span of incidence numbers, at a 95% confidence level. The percentage of the population above the age of 65, as well as that below the poverty line, are additional indicators with a significant impact on the case incidence rate, along with their interaction term.                                                                                                                                                                                                                                  |

|    |                             |                                                                                                                                                                            |         |            |                          |                                                                                                              |                     |                                                                                                   |                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|----|-----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------|--------------------------|--------------------------------------------------------------------------------------------------------------|---------------------|---------------------------------------------------------------------------------------------------|----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 18 | Fielding-Miller et al. [57] | Social determinants of COVID-19 mortality at the county level                                                                                                              | USA     | 08.05.2020 | Ecological study         | 2,743 counties across 50 states                                                                              |                     | Poverty level, rate uninsured                                                                     | Mortality            | In urban counties (n=114), only population density was significantly associated with COVID-19 mortality ( $b = 0.21$ , $p < 0.001$ ). In non-urban counties (n=2,629), all hypothesized social determinants were significantly associated with higher levels of mortality. Percentage of uninsured individuals was associated with lower reported COVID-19 mortality ( $b = -0.36$ , $p = 0.001$ ).                                                                                                                                                                                                       |
| 19 | Guha et al. [45]            | Community and Socioeconomic Factors Associated with COVID-19 in the United States: Zip code level cross sectional analysis                                                 | USA     | 22.04.2020 | Ecological study         | Population of New York City, Chicago, Richmond county of Detroit, Kings County of Seattle, Miami-Dade County | Until 11.04.2020    | Ethnicity/race, household income, poverty level, median house value, education                    | Incidence, mortality | A multivariable linear regression model noted that 1% increase in the proportion of residents above the age of 65 years, proportion of African American residents, proportion of females, persons per household and population density of the zip code increased the proportion of positive cases by 0.77%, 0.23%, 1.64%, 1.83% and 0.46% respectively ( $P < 0.01$ ) with only population density remaining significant in zip codes with greater than median number of cases. In zips with greater than median number of deaths, no community/socio-economic factor contributed significantly to death. |
| 20 | Ho et al. [70]              | Modifiable and non-modifiable risk factors for COVID-19: results from UK Biobank                                                                                           | England | 02.05.2020 | Retrospective cohort     | 428,225 participants, 340 cases                                                                              | 16.03. - 14.04.2020 | Ethnicity/race, Townsend index of area deprivation                                                | Incidence            | Non-modifiable risk factors included older age (RR 1.10 per 5 years), male sex (RR 1.64), black ethnicity (RR 1.86) and socioeconomic deprivation (RR 1.26 per SD increase in Townsend Index).                                                                                                                                                                                                                                                                                                                                                                                                            |
| 21 | Khanijahania [82]           | County-Level Proportions of Black and Hispanic populations, and Socioeconomic Characteristics in Association with Confirmed COVID-19 Cases and Deaths in the United States | USA     | 05.06.2020 | Ecological study         | 3,142 US counties in 50 states                                                                               |                     | Ethnicity/race, household income, education, financial hardship, insurance coverage, unemployment | Incidence, mortality | US counties with a higher proportion of black population, and a higher proportion of adults with less than high school diploma had disproportionately higher COVID-19 cases and deaths ( $\beta > 0$ , $p < 0.05$ ). A higher proportion of the Hispanic population was associated with higher confirmed cases ( $\beta = 1.03$ , 95% CI= 0.57-1.5), and higher housing cost to household income ratio was associated with higher deaths ( $\beta = 3.74$ , 95% CI= 2.14-5.37).                                                                                                                           |
| 22 | Khawaja et al. [71]         | Associations with covid-19 hospitalisation amongst 406,793 adults: the UK Biobank prospective cohort study                                                                 | England | 11.05.2020 | Prospective cohort study | 406,793 participants, 605 cases                                                                              | 16.03. - 16.04.2020 | Ethnicity/race, residential deprivation                                                           | Hospitalisation      | Major independent risk factors for hospitalisation with covid-19 were male sex (odds ratio 1.52; 95% confidence interval 1.28 to 1.81; $P < 0.001$ ), South Asian ethnicity (2.02; 1.28 to 3.17; $P = 0.002$ ) or black ethnicity (3.09; 2.18 to 4.38; $P < 0.001$ ) compared to white ethnicity and greater residential deprivation (1.92 for most deprived quartile compared to least deprived quartile; 1.50 to 2.47; $P < 0.001$ ).                                                                                                                                                                   |

|    |                    |                                                                                                                                                 |         |            |                  |                                                            |                     |                                                                                                              |                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|----|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------|------------------|------------------------------------------------------------|---------------------|--------------------------------------------------------------------------------------------------------------|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 23 | Li et al. [46]     | Multivariate Analysis of Factors Affecting COVID-19 Case and Death Rate in U.S. Counties: The Significant Effects of Black Race and Temperature | USA     | 24.04.2020 | Ecological study | 661 counties                                               | Until 14.04.2020    | Ethnicity, poverty level, violence rate                                                                      | Incidence, mortality          | Risk factors associated with increased cases and/or deaths per 100,000 included increased GDP per capita, increased percent Black, decreased percent Hispanic, decreased percent Asian, increased poverty and decreased access to primary care. Multivariate regression analyses demonstrated Black race is a risk factor for worse COVID-19 outcome independent of comorbidities, poverty, access to health care, and other mitigating factors.                                                                                                                                                                                                                                                                        |
| 24 | Liu et al. [72]    | Time courses of COVID-19 infection and local variation in socioeconomic and health disparities in England                                       | England | 29.05.2020 | Ecological study | Population of England                                      | 30.01. - 06.05.2020 | Ethnicity, Index of Multiple Deprivation                                                                     | Incidence                     | The higher case trajectory cluster (79 local authorities) had higher proportion of Black and Asian residents ( $p=0.03$ ; 29 $p=0.02$ ), higher multiple deprivation scores ( $p<0.001$ ). Local authorities with higher proportions of Black residents were more likely to belong to the high case trajectory cluster, even after adjusting for population density, deprivation, proportion of older adults and preventable mortality ( $p=0.04$ ).                                                                                                                                                                                                                                                                    |
| 25 | Maroko et al. [63] | Covid-19 and Inequity: A comparative spatial analysis of New York City and Chicago hot spots                                                    | USA     | 24.04.2020 | Ecological study | Population of New York City and Chicago                    | Until 13.04.2020    | Household density, ethnicity/race, education, household income, poverty level, unemployment rate, occupation | Incidence                     | In both cities, cold spots (clusters of low SARS-CoV-2 rate) demonstrated typical protective factors associated with the social determinants of health and the ability to social distance. These neighborhoods tended to be wealthier, have higher educational attainment, higher proportions of non-Hispanic white residents, and more workers in managerial occupations. Hot spots (clusters of high SARS-CoV-2 rate) also had similarities, such as lower rates of college graduates and higher proportions of people of color. It also appears to be larger households (more people per household), rather than overall population density, that may to be a more strongly associated with hot spots.               |
| 26 | Mukherji [47]      | The Social and Economic Factors Underlying the Impact of COVID-19 Cases and Deaths in US Counties                                               | USA     | 19.05.2020 | Ecological study | Population of 771 counties                                 | 30.03. - 18.04.2020 | Regional income, income inequality, poverty level, unemployment rate                                         | Incidence, mortality          | The paper finds that counties with high per capita personal income have high incidence of both reported cases and deaths. The unemployment rate is negative for deaths implying that places with low unemployment rates or higher economic activity have higher reported deaths. Counties with higher income inequality as measured by the Gini coefficient experienced more deaths and reported more cases. Counties with high concentrations of non-Hispanic Blacks, Native Americans, and immigrant populations have higher incidence of both cases and deaths.                                                                                                                                                      |
| 27 | Nayak et al. [73]  | Impact of Social Vulnerability on COVID-19 Incidence and Outcomes in the United States                                                          | USA     | 17.04.2020 | Ecological study | Population of 433 counties, 283,256 cases and 6,644 deaths | Until 04.04.2020    | Social Vulnerability Index (SVI)                                                                             | Incidence, case-fatality-rate | Higher SVI, indicative of greater social vulnerability, was associated with higher CFR (RR: 1.19 [1.05, 1.34], $p=0.005$ , per-1 unit increase), an association that strengthened after adjustment for age>65 years and comorbidities (RR: 1.63 [1.38, 1.91], $p<0.001$ ), and was further confirmed in a sensitivity analysis limited to six states with the highest testing levels. Although the association between overall SVI and COVID-19 incidence was not significant, the SVI sub-components of socioeconomic status and minority status were both predictors of higher incidence and CFR. A combination of high SVI ( $\geq 0.46$ ) and high adjusted CFR ( $\geq 2.3\%$ ) was observed in 28.9% of counties. |

|    |                         |                                                                                                                                               |                   |            |                          |                                               |                                      |                                                             |                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----|-------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|-------------------|------------|--------------------------|-----------------------------------------------|--------------------------------------|-------------------------------------------------------------|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 28 | Nazroo et al. [74]      | Evidence for ethnic inequalities in mortality related to COVID-19 infections: Findings from an ecological analysis of England and Wales       | England and Wales | 09.06.2020 | Ecological study         | Population of England and Wales               | Until 24.04.2020                     | Index of Multiple Deprivation, ethnicity                    | Mortality       | For every 1% rise in proportion of the population who are ethnic minority, COVID-19 related deaths increased by 5-10 (3-99 to 6-21) per million. This rise is present for each ethnic minority category examined. The size of this increase is a little reduced in a fully adjusted model, suggesting that some of the association results from ethnic minority people living in more densely populated, more polluted and more deprived areas.                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| 29 | Patel et al. [75]       | Race, Socioeconomic Deprivation, and Hospitalization for COVID-19 in English participants of a National Biobank                               | England           | 02.05.2020 | Prospective cohort study | 418,794 participants                          |                                      | Ethnicity, Townsend Index of Area Deprivation               | Hospitalisation | COVID-19 hospitalization was noted in 32 of 7,714 (0.4%) black participants, 28 of 10,614 (0.2%) Asian participants, and 489 of 400,438 (0.1%) white participants, with results largely consistent across English regions. In a logistic regression model adjusted for age, sex, and geographic region, both black participants (odds ratio 3.4; 95%CI 2.4-4.9) and Asian participants (odds ratio 2.1; 95%CI 1.5-3.2) were at increased risk as compared to white participants. The relationship between race and COVID-19 hospitalization was only modestly attenuated in a logistic regression model that additionally adjusted for Townsend Deprivation Index and household income--odds ratios for black and Asian participants of 3.1 (95%CI 2.0- 4.8) and 2.0 (95%CI 1.2-3.1) as compared to white participants respectively.                                                           |
| 30 | Plümper & Neumayer [48] | The COVID -19 Pandemic Predominantly Hits Poor Neighborhoods, or does it? Evidence from Germany                                               | Germany           | 18.05.2020 | Ecological study         | German population                             | Until 13.04. and 14.04. - 17.05.2020 | Regional income, education, rate of social welfare benefits | Incidence       | First time period: Higher incidence in regions with high income, high education, low rate of social welfare, in second time period high income and education are associated with lower incidence rates.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| 31 | Prats-Urbe et al. [76]  | Ethnicity, comorbidity, socioeconomic status, and their associations with COVID-19 infection in England: a cohort analysis of UK Biobank data | England           | 08.05.2020 | Retrospective cohort     | 415,582 participants, 1,416 tested, 651 cases | Until 14.04.2020                     | Ethnicity, Index of Multiple Deprivation                    | Incidence       | The incidence of COVID-19 was 0.61% (95% CI: 0.46%-0.82%) in Black/Black British participants, 0.32% (0.19%-0.56%) in 'other' ethnicities, 0.32% (0.23%-0.47%) in Asian/Asian British, 0.30% (0.11%-0.80%) in Chinese, 0.16% (0.06%-0.41%) in mixed, and 0.14% (0.13%-0.15%) in White. Compared with White participants, Black/Black British participants had an adjusted relative risk (RR) of 3.30 (2.39-4.55), Chinese participants 3.00 (1.11-8.06), Asian/Asian British participants 2.04 (1.36-3.07), 'other' ethnicities 1.93 (1.08-3.45), and mixed ethnicities 1.07 (0.40-2.86). Socioeconomic status (adjusted RR 1.93 (1.51-2.46) for the most deprived), obesity (adjusted RR 1.04 (1.02-1.05) per kg/m <sup>2</sup> ) and comorbid hypertension, chronic obstructive pulmonary disease, asthma, and specific renal diseases were also associated with increased risk of COVID-19. |

|    |                             |                                                                                                                                                                                                              |                |            |                            |                                                                   |                     |                                                        |                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|------------|----------------------------|-------------------------------------------------------------------|---------------------|--------------------------------------------------------|----------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 32 | Raisi-Estabragh et al. [19] | Greater risk of severe COVID-19 in non-White ethnicities is not explained by cardiometabolic, socioeconomic, or behavioural factors, or by 25(OH)-vitamin D status: study of 1,326 cases from the UK Biobank | United Kingdom | 02.06.2020 | Retrospective cohort study | 497,996 participants, 4,510 tested, 1,326 cases                   | 16.03. - 18.05.2020 | Ethnicity/race, Index of Multiple Deprivation, housing | Incidence            | There was over-representation of men and non-White ethnicities in the COVID-19 positive group. Non-White ethnicity, Townsend deprivation score and household overcrowding were independently associated with significantly greater odds of COVID-19. The pattern of association was consistent for men and women; socio-demographic and behavioural factors did not attenuate sex/ethnicity associations.                                                                                                                                                                                                                                                                                                                      |
| 33 | Rose et al. [58]            | Inequalities in COVID19 mortality related to ethnicity and socioeconomic deprivation                                                                                                                         | England        | 30.04.2020 | Ecological study           | Population of England                                             | Until 22.04.2020    | Ethnicity/race, poverty level                          | Mortality            | Local authorities with a greater proportion of residents from ethnic minority backgrounds had statistically significantly higher COVID-19 mortality rates, as did local authorities with a greater proportion of residents experiencing deprivation relating to low income. After adjusting for income deprivation and other covariates, each percentage point increase in the proportion of the population from BAME backgrounds was associated with a 1% increase in the COVID19 mortality rate [IRR=1.01, 95%CI 1.01–1.02]. Each percentage point increase in the proportion of the population experiencing income deprivation was associated with a 2% increase in the COVID19 mortality rate [IRR=1.02, 95%CI 1.01–1.04]. |
| 34 | Sy et al. [49]              | Socioeconomic disparities in subway use and COVID-19 outcomes in New York City                                                                                                                               | USA            | 30.05.2020 | Ecological study           | Population of New York City                                       | Until 26.04.2020    | Regional income, education, ethnicity, rate uninsured  | Incidence            | Increased subway use was associated with a higher rate of COVID-19 cases per 100,000 population when adjusted for testing effort (aRR=1.11; 95% CI: 1.03 - 1.19), but this association was weaker once we adjusted for median income (aRR=1.06; 95% CI: 1.00 - 1.12). All sociodemographic variables were significantly associated with the rate of positive cases per 100,000 population when adjusting for testing effort (except percent uninsured) and adjusting for both income and testing effort. The risk factor with the strongest association with COVID-19 was the percent of individuals in essential work (aRR = 1.59, 95% CI: 1.36 - 1.86).                                                                      |
| 35 | Takagi et al. [50]          | Ethnics and economics in COVID-19: Meta-regression of data from countries in the New York City metropolitan area                                                                                             | USA            | 24.05.2020 | Ecological study           | Population of 31 countries in the New York City Metropolitan Area | Until 20.05.2020    | Ethnicity/race, poverty level, household income        | Incidence, mortality | A slope (coefficient) of the univariable meta-regression line for COVID-19 prevalence was not significant for household income ( $P = .639$ ), whereas the coefficient was significantly positive for black (coefficient, 0.021; $P = .015$ ), Hispanic/Latino (0.033; $P < .001$ ), and poverty (0.039; $P = .02$ ), which indicated that COVID-19 prevalence increased significantly as black, Hispanic/Latino, and poverty increased. The multivariable model revealed that the slope was significantly positive for only Hispanic/Latino ( $P < .001$ ). The coefficient in the univariable model for COVID-19 fatality, however, was not significant for all the covariate.                                               |

|    |                             |                                                                                                                                                |         |            |                      |                                           |                     |                                                                                                        |                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|----|-----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|---------|------------|----------------------|-------------------------------------------|---------------------|--------------------------------------------------------------------------------------------------------|----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 36 | Takagi et al. [59]          | Meta-regression of COVID-19 prevalence/fatality on socioeconomic characteristics of data from top 50 U.S. large cities                         | USA     | 27.05.2020 | Ecological study     | Population of 50 US cities                | Until 22.05.2020    | Ethnicity/race, education, household income, insurance rate, poverty level, unemployment rate          | Incidence, mortality | COVID-19 prevalence was significantly negative for male sex, education attainment, computer and Internet use, and private health insurance. Whereas, the coefficient was significantly positive for black race, never matrimony, unemployment, and poverty. In the multivariable model, the coefficient was significantly negative for male sex ( $P = 0.036$ ) and computer use ( $P = 0.024$ ), and significantly positive for never matrimony ( $P < 0.001$ ). A coefficient for COVID-19 fatality was significantly negative for no health insurance, and significantly positive for elderly, unemployment and public coverage.                                                                                                                                                    |
| 37 | Vahidy et al. [51]          | Racial and Ethnic Disparities in SARS-CoV-2 Pandemic: Analysis of a COVID-19 Observational Registry for a Diverse U.S. Metropolitan Population | USA     | 28.04.2020 | Retrospective cohort | 4,513 participants, 754 cases             | 05.03. - 12.04.2020 | Ethnicity/race, household income                                                                       | Incidence            | Among 4,513 tested individuals, 754 (16.7%) tested positive. Overall mean (SD) age was 50.6 (18.9) years, 62% females and 26% were African American. African American race was associated with higher comorbidity burden, lower socio-economic status, and higher population density residence. In the fully adjusted model, African American race (vs. White; aOR, CI: 1.84, 1.49–2.27) and Hispanic ethnicity (vs. non-Hispanic; aOR, CI: 1.70, 1.35–2.14) had a higher likelihood of infection.                                                                                                                                                                                                                                                                                     |
| 38 | Whittle & Diaz-Artiles [52] | An ecological study of socioeconomic predictors in detection of COVID-19 cases across neighborhoods in New York City                           | USA     | 22.04.2020 | Ecological study     | Population of New York City, 64,955 cases | Until 05.04.2020    | Regional household income, unemployment rate, poverty level                                            | Incidence            | A decrease of \$10,000 median household income is associated with a 2.5% (95% CI: 1.0% to 4.1% $p = 0.002$ ) increase in detected COVID-19 cases.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| 39 | Williamson et al. [77]      | OpenSAFELY: factors associated with COVID-19-related hospital death in the linked electronic health records of 17 million adult NHS patients   | England | 07.05.2020 | Retrospective cohort | 17,425,445 people, 5,683 deaths           | 01.02. - 25.04.2020 | Index of Multiple Deprivation, ethnicity                                                               | Mortality            | Death from COVID-19 was strongly associated with being male (HR 1.99 (1.88-2.10)) older age and deprivation and comorbidities. Black people were at higher risk of death with only partial attenuation in hazard ratios from the fully adjusted model (age-sex adjusted HR 2.17 (1.84 - 2.57), fully adjusted HR 1.71 (1.44 - 2.02) with similar findings for Asian people (age-adjusted HR 1.95 (1.73-2.18), fully adjusted HR 1.62 (1.43-1.82).                                                                                                                                                                                                                                                                                                                                      |
| 40 | Xie & Li [64]               | Health and Demographic Impact on COVID-19 Infection and Mortality in US Counties                                                               | USA     | 11.05.2020 | Ecological study     | US population                             | 21.01. - 23.04.2020 | Regional poverty level, education, ethnicity/race, unemployment, household income, housing cost burden | Incidence            | The higher percentage of adults with a high school or less education, the higher of COVID-19 infection rate in log scale ( $b = 0.02113$ , 95% CI: 0.01491, 0.02734). With the higher housing cost burden, the COVID-19 infection rate in log scale significantly increased ( $b = 0.03451$ , 95% CI: 0.01786, 0.05116). With the higher percentage of Black or American Indian & Alaska Native, the COVID-19 infection rate in log scale increased significantly. With the higher of segregation index, the lower of COVID-19 infection rate in log scale ( $b = -0.00457$ , 95% CI: -0.00723, -0.00192). The percentage of Hispanic in the population showed a negative correlation with the infection rate of COVID-19 in log scale ( $b = -0.01236$ , 95% CI: -0.01957, -0.00515). |

|                                |    |                            |                                                                                                                                                                                            |                |            |                  |                                           |                       |                                                 |                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------|----|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|------------|------------------|-------------------------------------------|-----------------------|-------------------------------------------------|--------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reports and other publications | 41 | Chen & Krieger [60]        | Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county vs. ZIP code analyses                                                                | USA            | 21.04.2020 | Ecological study | Population of the USA, Illinois, New York | Data until 16.04.2020 | Area-based socioeconomic measures (ABSM)        | Incidence, mortality on county level | Highest death rates were observed in counties with the highest poverty; highest incidence in zip codes with highest ABSM.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|                                | 42 | Chen et al. [61]           | COVID-19 and the unequal surge in mortality rates in Massachusetts, by city/town and ZIP Code measures of poverty, household crowding, race/ethnicity, and racialized economic segregation | USA            | 09.05.2020 | Ecological study | Population of Massachusetts               | 01.01. - 15.04.2020   | Area-based socioeconomic measures (ABSM)        | Excess death rate linked to COVID-19 | Surge in excess death rates, both relative and absolute, was evident starting in early April, and was greater in city/towns and ZCTAs with higher poverty, higher household crowding, higher percentage of populations of color, and higher racialized economic segregation.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|                                | 43 | Public Health England [78] | Disparities in the risk and outcomes of COVID-19                                                                                                                                           | United Kingdom | 02.06.2020 | Ecological study | Population of England and Wales           |                       | Index of Multiple Deprivation (IMD), occupation |                                      | The trend in the number of diagnosed cases by deprivation quintile shows that cases in the least deprived group peaked earlier and lower than other groups and at 13 May, the cumulative number of cases and diagnosis rate was highest in the most deprived death. The trend in the number of diagnosed cases by deprivation quintile shows that cases in the least deprived group peaked earlier and lower than other groups and at 13 May, the cumulative number of cases and diagnosis rate was highest in the most deprived quintile. For three occupations the relative increase in deaths in 2020 was significantly higher than the average of 1.5: Caring Personal Services, Elementary Security Occupations, and Road Transport Drivers. Of these groups, the biggest increase was for Elementary Security Occupations, where deaths were 2.3 times higher in 2020 than in the same period in 2014 to 2018. Within these groups, there were three occupational 'unit groups' where the increase in deaths in 2020 was significantly higher than the increase for everyone aged 20 to 64. These were nursing auxiliaries and assistants, security guards and related occupations, and taxi and cab drivers and chauffeurs. The largest absolute increase was for workers in Caring Personal Services. There were 760 deaths from all causes among these workers in the period 21 March to 8 May 2020 for people aged 20 to 64. This is 346 more than in the same period in 2014 to 2018 and 74% had COVID-19 recorded as a cause of death. For workers in Construction and Building Trades, the number of deaths related to COVID-19 was slightly higher than the number of excess deaths. This indicates that deaths from other causes have gone down which may be due to a reduced risk of occupational related injuries over this time period. |

|    |                                                      |                                                                                                                             |                                     |            |                                              |                                                                    |                     |                                     |                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|----|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|-------------------------------------|------------|----------------------------------------------|--------------------------------------------------------------------|---------------------|-------------------------------------|---------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 44 | Office for National Statistics [79]                  | Deaths involving COVID-19 by local area and socioeconomic deprivation: deaths occurring between 1 March and 31 May 2020     | England and Wales                   | 12.06.2020 | Ecological study                             | 46,687 deaths linked to COVID-19                                   | 01.03. - 31.05.2020 | Index of Multiple Deprivation (IMD) | Mortality                                   | Looking at deaths involving the coronavirus (COVID-19), in England, the rate for the least deprived area (Decile 10) was 58.8 deaths per 100,000 population and the rate in the most deprived area (Decile 1) was 128.3 deaths per 100,000 population; this is 118% higher than the least deprived area. In the least deprived area, the age-standardised mortality rate for all deaths was 242.6 deaths per 100,000 population. In the most deprived area, the age-standardised mortality rate for all deaths was 92.2% higher than that of the least deprived, at 466.2 deaths per 100,000 population. The most deprived fifth of areas (quintile) in Wales had a rate of 109.5 deaths involving the coronavirus (COVID-19) per 100,000 population; this was nearly twice as high as the least deprived areas (57.5 deaths per 100,000 population) and over twice as high as the lowest mortality rate in Quintile 4 (50.5 deaths per 100,000 population). |
| 45 | Office for National Statistics [84]                  | Coronavirus (COVID-19) related deaths by occupation, England and Wales: deaths registered up to and including 20 April 2020 | England and Wales                   | 11.05.2020 | Analysis of official statistics              | 2,494 deaths in the working-age population of England and Wales    | Until 20.04.2020    | Occupation                          | Mortality                                   | Compared with the rate among people of the same sex and age in England and Wales, men working in the lowest skilled occupations had the highest rate of death involving COVID-19, with 21.4 deaths per 100,000 males (225 deaths); men working as security guards had one of the highest rates, with 45.7 deaths per 100,000 (63 deaths). Men and women working in social care, a group including care workers and home carers, both had significantly raised rates of death involving COVID-19, with rates of 23.4 deaths per 100,000 males (45 deaths) and 9.6 deaths per 100,000 females (86 deaths). Among men, a number of other specific occupations were found to have raised rates of death involving COVID-19, including: taxi drivers and chauffeurs (36.4 deaths per 100,000); bus and coach drivers (26.4 deaths per 100,000); chefs (35.9 deaths per 100,000); and sales and retail assistants (19.8 deaths per 100,000).                       |
| 46 | Intensive Care National Audit & Research Centre [80] | ICNARC report on COVID-19 in critical care 12 June 2020                                                                     | England, Wales and Northern Ireland | 12.06.2020 | Analysis of registry data / Ecological study | 9,777 patients from 289 ICU in England, Wales and Northern Ireland | Until 11.06.2020    | Index of Multiple Deprivation (IMD) | Admission to Intensive Care due to COVID-19 | 24,4 % of all COVID-19 ICU patients came from localities (postcode) in 4th Quintile and 25,3% from 5th Quintile of the distribution of the IMD.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |

ABSM = area-based socioeconomic measures, aOR = adjusted odds ratio, aRR = absolute risk reduction, BAME = black and minority ethnic, CI = confidence interval, GDP = gross domestic product, HR = hazard ratio, ICU = intensive care unit, IMD = Index of Multiple Deprivation, IRR = incident rate ratio, MRR = median rate ratio, NHS = National Health Service, OR = odds ratio, RR = relative risk, SD = standard deviation, SVI = Social Vulnerability Index, ZCTA = ZIP Code Tabulation Area